



BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Assessing the factors associated with mortality among patients recruited to the PLACID Trial (A Phase II, Open Label, Randomized Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease)

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-050571
Article Type:	Original research
Date Submitted by the Author:	23-Feb-2021
Complete List of Authors:	<p>Mammen, Joy; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Kumar, Snehil; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Thomas, Lovely; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Kumar, Gunjan ; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Zachariah, Anand; Christian Medical College and Hospital Vellore, Medicine</p> <p>Jeyaseelan, Lakshmanan; MBRU College of Medicine, College of Medicine; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Peter, John Victor; Christian Medical College, Vellore, Medical Intensive Care Unit</p> <p>Agarwal, Anup; ICMR, Clinical Trial and Health Systems Research Unit</p> <p>Mukherjee, Aparna; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Chatterjee, Pranab; ICMR, Translational Global Health Policy and Research Cell</p> <p>Bhatnagar, Tarun ; National Institute of Epidemiology, ICMR School of Public Health</p> <p>Rasalam, Jess; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Chacko, Binila; Christian Medical College, Vellore, Medical Intensive Care Unit</p> <p>Mani, Thenmozhi; Christian Medical College Vellore, Biostatistics</p> <p>Joy, Melvin; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Rupali, Priscilla; Christian Medical College Vellore, Infectious Diseases</p> <p>Murugesan, Malathi; Christian Medical College Vellore, Hospital Infection Control Committee</p> <p>Daniel, Dolly; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Latha, B; Madras Medical College, Transfusion Medicine</p> <p>Bundas, Sunita; SMS Medical College and Hospital, Transfusion Medicine</p> <p>Kumar, Vivek; Sir HN Reliance Foundation Hospital and Research Centre, Critical Care</p>

	Dosi, Ravi; Sri Aurobindo Institute of Medical Sciences, Respiratory Medicine
	Khambholja, Janakkumar; Smt NHL Municipal Medical College, Internal Medicine
	de Souza, Rosemarie; BYL Nair Charitable Hospital, Department of Medicine
	Chander, B; Gandhi Medical College and Hospital, Internal Medicine
	Bahadur, Shalini; Government Institute of Medical Sciences, Pathology
	Dube, Simmi; Gandhi Medical College Bhopal, Internal Medicine
	Suri, Amit; Atal Bihari Vajpayee Institute of Medical Sciences and Ram Manohar Lohia Hospital, Pulmonary Medicine
	Jindal, Aikaj ; Satguru Pratap Singh Hospitals, Transfusion Medicine
	Shrivastav, Om ; Kasturba Hospital for Infectious Diseases, Infectious diseases
	Barge, Vijay; RSCM Government Medical College, Medicine
	Bajpayee, Archana; AIIMS Jodhpur, Transfusion Medicine
	Malhotra, Pankaj; Post Graduate Institute of Medical Education and Research, Department of Internal Medicine
	Singh, Neha; AIIMS Patna, Transfusion Medicine
	Tambe, Muralidhar; Department of Community Medicine, B J Govt. Medical College
	Sharma, Nimisha; ESIC Medical College and Hospital Faridabad, Transfusion Medicine
	Bhat, Shreepad; Smt Kashibai Navale Medical College and General Hospital, Internal Medicine
	Kaulgud, Ram; Karnataka Institute of Medical Sciences, Internal Medicine
	Gurtoo, Anil; Lady Hardinge Medical College, Internal Medicine
	Reddy, Himanshu; King George Medical College, Internal Medicine
	Upadhyay, Kamlesh; Byramjee Jeejabhoy Medical College, Internal medicine
	Jain, Ashish ; Mahatma Gandhi Medical College and Hospital, Respiratory Medicine
	Patel, Tinkal; Government Medical College, Internal Medicine
	Nagori, Irfan; GMERS Medical College Gotri Vadodara, Medicine
	Jha, Pramod ; Sumandeep Vidyapeeth University, Internal Medicine
	Babu, KV; Sri Venkateswara Institute of Medical Sciences, Transfusion Medicine
	Panjwani, Sunil; Government Medical College, Internal Medicine
	Aparna, C; Kurnool Medical College
	Natarajan, M; Madurai Medical College, Internal Medicine
	Baldi, Milind; Mahatma Gandhi Memorial Medical College, Internal Medicine
	Khadke, Vrushali; Poona Hospital and Research Centre, Interventional Pulmonology
	Dua, Seema; Super Speciality Paediatric Hospital and Teaching Hospital, Transfusion Medicine
	Singh, Ravindraa; Aditya Birla Memorial Hospital, Transfusion Medicine
	Sharma, Ashish; R. D. Gardi Medical College, Medicine
	Sharma, Jayashree; Seth GS Medical College and KEM Hospital, Transfusion Medicine
	Gokhale, Yojana; Lokmanya Tilak Municipal Medical College and General Hospital, Internal Medicine
	Yadav, Pragya; National Institute of Virology, Maximum Containment Laboratory, ICMR
	Sapkal, Gajanan; National Institute of Virology, Diagnostic Virology Group, ICMR
	Kaushal, Himanshu; National Institute of Virology, Human Influenza Group, ICMR
	Kumar, Saravana; National Institute of Epidemiology, Epidemiology and Biostatistics Division

Keywords:

COVID-19, Public health < INFECTIOUS DISEASES, Adult intensive & critical care < INTENSIVE & CRITICAL CARE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

PLACID TRIAL Mortality Assessment**Assessing the factors associated with mortality among patients recruited to the PLACID Trial
(A Phase II, Open Label, Randomized Controlled Trial to Assess the Safety and Efficacy of
Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease)**

Joy J Mammen, Snehil Kumar, Lovely Thomas, Gunjan Kumar, Anand Zachariah, Lakshmanan
Jeyaseelan, John Victor Peter, Anup Agarwal, Aparna Mukherjee, Pranab Chatterjee, Tarun
Bhatnagar, Jess Elizabeth Rasalam, Binila Chacko, Thenmozhi Mani, Melvin Joy, Priscilla Rupali,
Malathi Murugesan, Dolly Daniel, B Latha , Sunita Bundas, Vivek Kumar, Ravi Dosi, Janakkumar R
Khambholja, Rosemarie de Souza, B Thrilok Chander, Shalini Bahadur, Simmi Dube, Amit Suri, Aikaj
Jindal, Om Shrivastav, Vijay Barge, Archana Bajpayee, Pankaj Malhotra, Neha Singh, Muralidhar
Tambe, Nimisha Sharma, Shreepad Bhat, Ram S Kaulgud, Anil Gurtoo, D Himanshu Reddy, Kamlesh
Upadhyay, Ashish Jain, Tinkal C Patel, Irfan Nagori, Pramod R Jha, KV Sreedhar Babu, C Aparna, Sunil
Jodharam Panjwani, M Natarajan, Milind Baldi, Vrushali Khirid Khadke, Seema Dua, Ravindraa
Singh, Ashish Sharma, Jayashree Sharma, Yojana A Gokhale, Pragya D Yadav, Gajanan Sapkal,
Himanshu Kaushal, V Saravana Kumar

Corresponding Author:**Joy J Mammen MD**

Professor & Head

Address

Department of Transfusion Medicine and Immunohaematology

Christian Medical College Vellore

Tamil Nadu, India

Pincode 632004

E mail joymammen@cmcvellore.ac.in**Phone** +91 416 228 2536**Fax** +91 416 223 2035

PLACID TRIAL Mortality Assessment

Abstract

Objective:

Large data on the clinical characteristics and outcome of COVID-19 in the Indian population is scarce. We analyzed the factors associated with mortality in a cohort of moderately ill COVID-19 patients enrolled in a randomized trial on convalescent plasma.

Setting:

39 public and private hospitals across India.

Participants:

Of the 464 patients recruited, two were lost to follow-up, nine withdrew consent and two patients did not receive the intervention after randomization. The cohort of 451 participants with known outcome at 28-days were analyzed.

Primary outcome measure:

Factors associated with all-cause mortality at 28 days post-enrolment.

Results:

The mean (SD) age of was 51±12.4 years; 76.7% were male. Admission SOFA score was 2.4±1.1. Non-invasive ventilation, invasive ventilation and vasopressor therapy were required in 98.9%, 8.4% and 4.0% respectively. The 28-day mortality was 14.4%. Median time from symptom onset to hospital admission was similar (p=1.0) in survivors (4 days; IQR 3-7) and non survivors (4 days; IQR 3-6). Patients with two or more co-morbidities had 2.25 (95%CI:1.17–4.32, p=0.014) times risk of death. When compared with survivors, admission IL-6 levels were higher (p<0.001) in non-survivors and increased further on Day 3. On multivariable regression analysis, severity of illness (HR 1.21, 95%CI:1.07-1.36, p=0.002), PaO₂/FiO₂ ratio <100 (3.37, 1.54-7.41, p=0.002), Neutrophil Lymphocyte ratio (NLR) >10 (9.38, 3.67-24.0, p<0.001), D-dimer >1.0mg/l (2.51,1.14-5.51, p=0.022), ferritin >500ng/ml (2.66, 1.46-4.85, p=0.001) and LDH ≥450 IU/L (2.96, 1.61-5.45, p=0.001) were significantly associated with death.

Conclusion:

In this cohort of moderately ill COVID-19 patients, severity of illness, underlying co-morbidities and higher levels of inflammatory markers were significantly associated with death.

Trial Registration:

The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775).

PLACID TRIAL Mortality Assessment

Article Summary

Strengths and limitations of this study

Strengths

There is no study from India, with representation from multiple states that has detailed the clinical profile and evaluated for factors associated with death. This study may help with strategic planning at a national level.

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the present analysis need not be adjusted for convalescent plasma intervention.

There may be variability of treatment provided in the multiple centres, however, care was taken that patients received best standard of care for covid-19 dictated by the best available evidence at the time and guidelines for the management of covid-19 issued by health authorities of the Indian government.

Limitations

The laboratory and biomarker assays for ferritin, lactate dehydrogenase, C reactive protein, and D-dimer were conducted using tests from different manufacturers.

Participants of this study may not comprise a true observational cohort as this was a post hoc analysis of a randomized control trial data and extrapolation to the general population must be carefully qualified.

PLACID TRIAL Mortality Assessment

Introduction:

The first human case of Corona Virus Disease 19 (COVID-19) caused by the novel coronavirus (named Severe acute respiratory syndrome coronavirus 2, SARS-CoV-2) was reported in Wuhan City, China in December 2019. On 30 January 2020, the World Health Organization (WHO) declared that the outbreak of COVID-19 constituted a Public Health Emergency of International Concern (1). Based on the high levels of global spread and the severity of COVID-19, on 11 March 2020, the Director-General of the WHO declared the COVID-19 outbreak a pandemic (2). The sudden outbreak followed by rapid spread in a globalized world, resulted in a major medical burden, besides affecting socio-economic wellbeing among all nations.

In India, the disease was first detected on 30 January 2020 in the state of Kerala, in a student who returned from Wuhan (3). After a brief, initial respite, the virus has spread at a rapid pace in India, resulting in more than 10 million confirmed cases as of December, 2020 with more than 145,000 deaths (4).

Patients diagnosed with COVID-19 have primarily respiratory symptoms. Most patients diagnosed with COVID-19 experience mild to moderate respiratory illness, fever, dry cough, fatigue and recover without requiring special treatment (5). Oxygen desaturation is the hallmark of progression. Patients with underlying medical problems like cardiovascular disease, diabetes, chronic respiratory disease, and cancer are more likely to develop serious illness. These patients may develop viral pneumonia, with resultant dyspnea and hypoxemia which may progress to respiratory or multi- system failure and even death (6). There is paucity of large-scale data of the clinical characteristics, outcomes of COVID-19 in the Indian population and evaluation of risk factors with an unfavorable outcome at a national level. Identification of such potential risk factors is important to anticipate medical treatment and to reduce the mortality burden for severe COVID-19 illness by proactive interventions.

The Indian Council of Medical Research (ICMR) conducted a randomized trial (A Phase II, Open Label, Randomized Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease, PLACID TRIAL) to determine the effectiveness and safety of convalescent plasma in patients with moderate COVID-19 to limit progression to severe disease (7). Patients enrolled received standard of care for COVID-19 in keeping with the institutional protocols, based on the best available evidence at the time and guidelines for the management of COVID-19 issued by national health authorities. Participants in the

PLACID TRIAL Mortality Assessment

intervention arm received two doses of 200 mL of convalescent plasma, transfused 24 hours apart, in addition to standard of care. The control arm did not receive any additional therapy. The study concluded that the use of convalescent plasma was not associated with a reduction in 28-day mortality (7).

The aim of this analysis was to identify risk factors associated with mortality by mining the data collected from the cohort enrolled in the PLACID TRIAL (7).

Methods

Participants

The study enrolled patients from 39 different hospitals, of which, 29 were teaching public hospitals and 10 were private facilities spread across 14 states and union territories. Patients over the age of 18 years who were confirmed to have COVID-19 based on a positive SARS-CoV-2 RT-PCR test and moderately ill with either a partial pressure of oxygen in arterial blood/fraction of inspired oxygen ($\text{PaO}_2/\text{FiO}_2$) ratio between 200-300 or respiratory rate $>24/\text{min}$ and decreased oxygen saturation on room air ($\text{SpO}_2 < 93\%$) were included. Patients were followed up for 28 days and assessed for their health status and all-cause mortality. Ethics approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals. Written consent was obtained from patients or their families before enrolling in the study.

Data

Data were obtained from the ICMR PLACID TRIAL database collected in structured paper case record forms and entered in Research Electronic Data Capture system (REDCap, version 8.5 Vanderbilt University, TN). The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775). After the trial was completed, based on cooperative agreement between the centers, and IRB permission, the data was shared and analyzed further, to explore for other meaningful results. No separate ethical clearance was taken for this study.

Demographic, clinical, laboratory tests and outcome data were collected prospectively. Clinical symptoms need for organ support (respiratory, hemodynamic), laboratory tests (complete blood count, coagulation profile, serum biochemical profile, renal and liver function tests) were monitored serially on day of enrollment (day 0) and on day 1, 3, 5, 7, 14 and 28. Inflammatory biomarkers

PLACID TRIAL Mortality Assessment

[lactate dehydrogenase (LDH), serum ferritin, and C-reactive protein (CRP)] and were tested at admission and on day 3 and 7 whereas, Interleukin 6 (IL-6) was done at admission and on day 3. The outcome of interest was all-cause day 28 mortality. We evaluated for association between laboratory parameters and mortality.

Statistical Methods:

Mean and standard deviation (SD) or median and inter-quartile range (IQR) were used for continuous variables as appropriate, and categorical variables number and proportions were used. To find the association between mortality and study variables, Chi-square test/Fisher’s exact test were used. To find the mean difference across the groups, independent t test was used. Similarly, Mann Whitney U test was used to compare median difference. Clinically important baseline variables and time dependent covariates were included in multivariable Cox proportional hazards model. Two multivariate models were developed. The first model included clinical and laboratory parameters tested on day 0, 1, 3, 5, 7, 14 and 28 while the second included inflammatory biomarkers tested on day 0, 3 and 7 after adjusting for age and comorbidities. Variables included parameters that were strongly associated with mortality at univariable analysis or known from previous literature to be strongly associated with outcome. The model assumption was verified using log-log S (t) plots and Global test. A p-value of less than 0.05 levels was considered as statistically significant. All statistical analyses were performed using STATA version 16.0 (StataCorp. 2019. College Station, TX).

Patient and public involvement

Patients and the public were not involved in the design, conduct, reporting, or dissemination plans of our research. The study results will be disseminated to the study participants via their treating doctors.

Results:

The PLACID Trial recruited 464 eligible patients for the study. The primary outcome at 28 days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomization and two patients did not receive the intervention after randomization as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients (*supplementary*).

PLACID TRIAL Mortality Assessment

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the analysis did not adjust for convalescent plasma intervention. The distribution of patients in intervention and control arms were 50.3% (n=227) and 49.7% (n=224), respectively. The mean (SD) age of the cohort was 51 ± 12.4 years; 76.7% were male. Table 1 shows distribution of demographic variables and clinical parameters in the study population.

The most common presenting symptoms were shortness of breath (91.6%), fatigue (78.7%), cough (68.5%) and fever (35%). Comorbidities were present in 59.9% of patients; 28.2% had two or more comorbidities. The most frequent comorbidities were diabetes (43.5%), hypertension (37.5%), obesity (6.9%) and Chronic Obstructive Pulmonary Disease (COPD) (3.3%). There was history of smoking in 8.2%. The time from onset of symptoms to admission was four days (IQR 3-7 days). Majority of the patients required non-invasive (98.9%) ventilatory support. The median duration of respiratory support was six days (IQR 4-10 days). In this cohort, 4% patients required vasopressor support. None of the patients required Extra Corporeal Membrane Oxygenation (ECMO) or dialysis support.

The all-cause mortality at 28 days was 14.4% (95%CI: 11.5-17.9, n=65). Median time from symptom onset to hospital admission was four days in survivors (IQR 3-7 days) and non survivors (IQR 3-6 days). The frequency of shortness of breath, cough and fatigue were similar in survivors and non survivors; however, the presence of fever at admission was significantly (p=0.04) associated with death (table 1). Other than COPD and CKD (chronic kidney disease), other comorbidities were not significantly associated with death (table 1). Admission SOFA score was higher in non survivors. The need for invasive mechanical ventilation, duration of invasive mechanical ventilation and vasopressor therapy were associated with death (table 1).

On univariable analysis (table 2), there was an association between increasing age and mortality. Patients with two or more comorbidities had a 2.25 (1.17-4.32, p=0.014) times increased chance of mortality. There was a strong mortality association for admission platelet count < 100 * 10⁹/L (HR 6.53, 95%CI: 3.10-13.75, p < 0.001), neutrophil lymphocyte ratio (NLR) > 10 (27.49, 11.67-64.75, p < 0.001), LDH ≥ 450 IU/L (4.88, 2.73-8.72, p < 0.001), D-dimer > 1mg/L (3.35, 1.55-7.23, p=0.002) and ferritin > 500ng/ml (4.09, 2.31-7.23, p < 0.001). Admission IL-6 levels were significantly (p < 0.001) higher (76.00, 18.27-171.77) in non survivors than in survivors (18.51, 4.26-56.86). By day 3, IL-6 levels dropped to 11.6 (2.64-45.84) in survivors while it nearly doubled in non-survivors

PLACID TRIAL Mortality Assessment

(140.35, 21.56-427.36). Univariate analysis of CRP did not show any statistical significance (1.0003, 0.99-1.001, p=0.150).

The need for invasive ventilation and vasopressors were associated with death (table 2). Increasing SOFA score was associated with mortality (1.61, 1.48-1.75, p <0.001). The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score progressively increased between the two groups over time (figure 1). Mortality proportionately also increased with lower PaO₂/FiO₂ values with hazard ratio of 23.11 (12.81-41.69, p <0.001) in severe group as compared to mild group.

Two models were run for multivariable Cox proportional hazards regression analysis over a period of time. Model A included age, comorbidities, PaO₂/FiO₂, NLR and SOFA score. Model A revealed significant hazard ratios for PaO₂/FiO₂ ratio < 100 (3.37, 1.54-7.41, p=0.002), NLR > 10 (9.38, 3.67-23.99, p <0.001), SOFA score (1.21, 1.07- 1.36, p=0.002) after adjusting for age and comorbidities. Model B included age, comorbidities, D-dimer, ferritin and LDH. D-dimer >1 mg/L (2.51, 1.14-5.51, p=0.022), ferritin >500 ng/ml (2.66, 1.46-4.85, p=0.001) and LDH ≥ 450 IU/L (2.96, 1.61-5.45, p=0.001), were associated with mortality after adjusting for age and comorbidities (table 2). IL-6 was omitted from the model as it was not measured on Day 7. CRP was not included in the model as it did not show significant difference between the two groups.

Discussion:

In this study that enrolled patients from across India, we were able to identify clinical, biomarkers (D-dimer, LDH, ferritin) and SOFA score as factors that could indicate increased risk of death in moderately ill COVID-19 patients from PLACID trial cohort. The definition of clinical grading of severity is different in India as compared to other countries (8-12). Mortality of patients critically ill with COVID-19 varies significantly among the published case series, ranging from 16% to 78% (13-19). Similarly two studies from Wuhan which included moderately as well as critically ill patients have shown mortality rate of 3.77% and 14.14% (20,21). This wide variability can be explained by differences in the age of the population, distribution of risk factors, health system response across different countries, varied treatment protocols and follow-up. In a series of critically ill patients in China, 28-day ICU mortality was 61.5% (22). In a multicentric study from Italy, the mortality risk for patient without respiratory failure at admission was of 1% after 15 days while survival in patients with a moderate-to-severe respiratory failure (PaO₂/FiO₂ ≤200 mm Hg) at admission was only 56% at 15 days (23). The fatality rate reported in Europe and the United States of America is significantly

PLACID TRIAL Mortality Assessment

higher than in China (24). Therefore, findings obtained in a specific country might not be automatically extrapolated and national cohorts must be studied.

In our study population, mortality increased with age, which is similar to the pattern observed in other countries affected by COVID-19. Age seems to affect the time from hospitalization to death. Age-specific death rates was quite similar in studies from Asia, Europe and North America (25). South Korea, Italy, France, Germany, England and Wales, and Spain COVID-19 attributed mortality rates rise by about 12%/year while the United States and Wuhan, China showed a slower rate of increase about 9.5%/year of age (26). In a meta-analysis of 61,11,583 subjects, 23.2% of patients were aged ≥ 80 years and showed an average mortality rate of 12.10%, the lowest being in China (3.1%) and the highest in the United Kingdom (20.8%) and New York State (20.99%). In the same study, highest mortality rate was observed in patients aged ≥ 80 years. The largest increase in mortality risk was observed in patients aged 60 to 69 years compared with those aged 50 to 59 years (odds ratio 3.13, 95% CI: 2.61-3.76) (27).

Presence of comorbidities significantly increases the death risk of COVID-19. A higher risk of mortality was seen in our patients who had CKD and COPD. Meta-analysis, including 1389 COVID-19 patients with 19.7% having severe disease showed a significant association of CKD with severe COVID-19 with pooled odds ratio as 3.03 (28). Similarly, the estimated mortality risk in patients with COPD was three times of those without ($p < 0.05$) (29). We found that 43.5% of our patients had diabetes which is markedly higher as compared to patients from Korea which showed that 16.97% had diabetes mellitus (30). Our analysis showed that the presence of diabetes was not significantly different between survivors and non survivors (42.5% vs. 49.2%, $p = 0.310$), in contrast to the study from South Korea (30) which showed a much higher mortality among diabetic patients than in those without (20.0% vs. 4.8%). Hypertension and obesity were not significantly different among survivors and non-survivors in our study. However, the presence of two or more comorbidities was associated with mortality in our study.

Univariable Cox proportional hazards regression modeling identified several other prognostic markers for mortality, most notably age ≥ 60 years, $\text{PaO}_2/\text{FiO}_2$ ratio < 100 , NLR > 10 , platelet count $< 100 \times 10^9/\text{L}$, ferritin $> 500\text{ng/ml}$, LDH $> 450 \text{ IU/L}$ and D-dimer $> 1\text{mg/L}$. Our study showed similar findings when compared with studies from Wuhan (31). Older age, leukocytosis, and high LDH level have been reported to be risk factors associated with in-hospital death in other studies also (32–34).

PLACID TRIAL Mortality Assessment

IL-6 levels were significantly different in survivors and non-survivors at admission. By day 3 survivors had reducing IL-6 while it nearly doubled in non survivors.

Mortality was higher among patients requiring invasive mechanical ventilation (HR 19.57, 11.81-32.41, $p<0.001$) and those requiring vasopressors (HR 11.36, 6.47-19.96, $p<0.001$). However, the median duration of invasive ventilation for survivors was 12 days (2, 14) and that for non-survivors was one day (1,3). These results suggest that patients with acute respiratory failure from COVID-19 may recover, even with severe disease requiring longer ventilation, and that probably the sickest patients die early reflecting lower duration of invasive ventilation in non survivors. Therefore, invasive ventilation should be timely and effectively provided.

In our study, the SOFA score was recognized as a valuable tool that could be used to prognosticate outcome of patients with COVID-19. Univariate and multivariate regression analysis both showed that the increase in SOFA score was related to mortality, with a clearly divergent pattern between the two groups. Thus, an increasing SOFA score over time may be a factor that can be used to identify a subset of patients who may have an unfavorable outcome. Studies have shown that the SOFA score could be used to evaluate severity and 60-day mortality of COVID-19 with the optimal cut-off score of 5 (35).

Limitations of the study includes the variability of treatment provided in the multiple centres. The participants of this study may not comprise a true observational cohort as this was a post hoc analysis of a randomized control trial data and extrapolation to the general population must be carefully qualified.

The risk factors identified in this study include older age with two or more comorbidities, mainly history of COPD and CKD. Multivariate analysis showed lymphopenia, lower PaO₂/ FiO₂ ratio, increased LDH, ferritin and D-dimer were significantly associated with mortality. A rising IL-6 may portend a poor prognosis. Serial SOFA score can be used for prognostication. Understanding the symptoms, burden of comorbidities, systematic monitoring key laboratory parameters offer new methods of improving reducing mortality in COVID-19.

PLACID TRIAL Mortality Assessment

FOOTNOTES:

Authors Contributions:

Study design: JJM, LJ, JVP

Clinical Management: SK, LT, AZ, JER, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG.

Data collection: JJM, SK, LT, AZ, AA, AM, GK, PC, TB, JER, PR, MM, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG, PDY, GS, HK, VSK.

Data Analysis: JJM, SK, LJ, TM, MJ, PR, MM, DD, JVP.

Data Interpretation: JJM, SK, AZ, LJ, JER, TM, MJ, DD, JVP

Manuscript writing: JJM, SK, LT, AZ, LJ, JER, BC, TM, MJ, PR, MM, DD, JVP, AA, AM, GK, HK, PC, TB

Study Administration: JJM, BC, JVP, AA, AM, GK, HK, PC, TB

Funding support: The funding source for the primary study was ICMR (RFC No. ECD/NTF/20/2020-21/Covid Dated 23.07.2020). However, they had no financial interest in the investigational product. No other author has any competing financial or non-financial interest.

Competing interests: All authors have completed the ICMJE uniform disclosure form at www.icmje.org. LJ was member of the independent Data and Safety Monitoring Board for the PLACID trial. He did not receive any remuneration for the primary study. This is a secondary study conducted using data collected from patients enrolled during the PLACID trial. No new patient was enrolled during this study.

Ethical Approval: Ethical approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) based in the National Center for Disease Informatics and Research, Indian Council of Medical Research, Bengaluru, Karnataka, as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals.

Data availability statement: Data will be made available, upon request, and must be accompanied by a brief proposal outlining the analysis plan. A signed data access agreement might be needed to ensure data safety and compliance with national rules about data sharing.

PLACID TRIAL Mortality Assessment

Acknowledgement: We acknowledge ICMR for providing us the data collected during PLACID trial which was analyzed in the present study.

References:

1. Mullen L, Potter C, Gostin LO, Cicero A, Nuzzo JB. An analysis of International Health Regulations Emergency Committees and Public Health Emergency of International Concern Designations. *BMJ Glob Health*. 2020 Jun 1;5(6):e002502.
2. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Bio-Medica Atenei Parm*. 2020 19;91(1):157–60.
3. Perappadan BS. India’s first coronavirus infection confirmed in Kerala. *The Hindu* [Internet]. 2020 Jan 30 [cited 2020 Nov 16]; Available from: <https://www.thehindu.com/news/national/indias-first-coronavirus-infection-confirmed-in-kerala/article30691004.ece>
4. MoHFW | Home [Internet]. [cited 2021 Jan 3]. Available from: <https://www.mohfw.gov.in/>
5. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol Orlando Fla*. 2020 Jun;215:108427.
6. Wu C, Chen X, Cai Y, Xia J ’an, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med*. 2020 01;180(7):934–43.
7. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre randomised controlled trial (PLACID Trial). *BMJ* [Internet]. 2020 Oct 22 [cited 2020 Nov 16];371. Available from: <https://www.bmj.com/content/371/bmj.m3939>
8. Varghese GM, John R, Manesh A, Karthik R, Abraham OC. Clinical management of COVID-19. *Indian J Med Res*. 2020 May 1;151(5):401.
9. Clinical management of COVID-19 [Internet]. [cited 2021 Jan 3]. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19>
10. Zhang S-Y, Lian J-S, Hu J-H, Zhang X-L, Lu Y-F, Cai H, et al. Clinical characteristics of different subtypes and risk factors for the severity of illness in patients with COVID-19 in Zhejiang, China. *Infect Dis Poverty*. 2020 Jul 8;9(1):85.
11. Xia L, Chen J, Friedemann T, Yang Z, Ling Y, Liu X, et al. The Course of Mild and Moderate COVID-19 Infections—The Unexpected Long-Lasting Challenge. *Open Forum Infect Dis* [Internet]. 2020 Sep 1;7(ofaa286). Available from: <https://doi.org/10.1093/ofid/ofaa286>

PLACID TRIAL Mortality Assessment

12. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020 Jun 1;20(6):669–77.
13. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al. Risk Factors Associated With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy. *JAMA Intern Med.* 2020 01;180(10):1345–55.
14. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet Lond Engl.* 2020 15;395(10223):497–506.
15. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, et al. Clinical Course and Outcomes of 344 Intensive Care Patients with COVID-19. *Am J Respir Crit Care Med.* 2020 Apr 8;201(11):1430–4.
16. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA.* 2020 Mar 17;323(11):1061–9.
17. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, et al. Covid-19 in Critically Ill Patients in the Seattle Region - Case Series. *N Engl J Med.* 2020 21;382(21):2012–22.
18. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA.* 2020 28;323(16):1612–4.
19. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet Lond Engl.* 2020 28;395(10229):1054–62.
20. Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect.* 2020 Jun 1;26(6):767–72.
21. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ.* 2020 Mar 26;368:m1091.
22. Yang X, Yu Y, Xu J, Shu H, Xia J 'an, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med.* 2020 May 1;8(5):475–81.
23. Santus P, Radovanovic D, Saderi L, Marino P, Cogliati C, De Filippis G, et al. Severity of respiratory failure at admission and in-hospital mortality in patients with COVID-19: a prospective observational multicentre study. *BMJ Open [Internet].* 2020 Oct 10;10(10). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7549463/>

PLACID TRIAL Mortality Assessment

24. Yamamoto N, Bauer G. Apparent difference in fatalities between Central Europe and East Asia due to SARS-COV-2 and COVID-19: Four hypotheses for possible explanation. *Med Hypotheses*. 2020 Nov;144:110160.

25. Demographics of COVID-19 Deaths [Internet]. Ined - Institut national d'études démographiques. [cited 2020 Nov 16]. Available from: <https://dc-covid.site.ined.fr/en/>

26. Goldstein JR, Lee RD. Demographic perspectives on the mortality of COVID-19 and other epidemics. *Proc Natl Acad Sci U S A*. 2020 Sep 8;117(36):22035–41.

27. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583 Subjects. *J Am Med Dir Assoc*. 2020 Jul 1;21(7):915–8.

28. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *Int Urol Nephrol*. 2020 Mar 28;1–2.

29. Venkata VS, Kiernan G. COVID-19 AND COPD: POOLED ANALYSIS OF OBSERVATIONAL STUDIES. *CHEST*. 2020 Oct 1;158(4):A2469.

30. D A, K L, Ds L, Ys L, Ss M. Mortality Rate and Predictors of Mortality in Hospitalized COVID-19 Patients with Diabetes. *Healthc Basel Switz* [Internet]. 2020 Sep 13 [cited 2020 Nov 16];8(3). Available from: <https://europepmc.org/article/med/32933191>

31. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* [Internet]. 2020 May 7;55(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144257/>

32. Li M, Cheng B, Zeng W, Chen S, Tu M, Wu M, et al. Analysis of the Risk Factors for Mortality in Adult COVID-19 Patients in Wuhan: A Multicenter Study. *Front Med* [Internet]. 2020 [cited 2020 Nov 16];7. Available from: <https://www.frontiersin.org/articles/10.3389/fmed.2020.00545/full>

33. Albitar O, Ballouze R, Ooi JP, Sheikh Ghadzi SM. Risk factors for mortality among COVID-19 patients. *Diabetes Res Clin Pract*. 2020 Aug 1;166:108293.

34. Holman N, Knighton P, Kar P, O'Keefe J, Curley M, Weaver A, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol*. 2020 Oct 1;8(10):823–33.

35. Zheng Yang, Qinming Hu, Fei Huang, Shouxin Xiong, Yi Sun. The prognostic value of the SOFA score in patients with COVID-19: a retrospective, observational study. 2020 Oct 26 [cited 2020 Nov 20]; Available from: <https://europepmc.org/article/ppr/ppr231113>

PLACID TRIAL Mortality Assessment

Table 1: Distribution of demographic variables and clinical parameters in enrolled patients and comparison of survivors and non-survivors in the cohort

Variables		Overall (n=451) N (%)	Survivor (n=386) N (%)	Non survivor (n=65) N (%)	P value
Age (Mean ± SD)		51±12.4	50±12.4	56±11.3	<0.001*
Age	≤ 40	104 (23.1)	97 (25.1)	7 (10.8)	0.004
	41-59	225 (49.9)	194 (50.3)	31 (47.7)	
	≥60	122 (27.0)	95 (24.6)	27 (41.5)	
Gender: Male		346 (76.7)	294 (76.2)	52 (80.0)	0.499
Blood group	A	104 (23.1)	91 (23.6)	13 (20.0)	0.530
	B	164 (36.4)	140 (36.3)	24 (36.9)	
	AB	25 (5.5)	19 (4.9)	6 (9.2)	
	O	158 (35.0)	136 (35.2)	22 (33.9)	
History of smoking		37 (8.2)	32 (8.3)	5 (7.7)	0.866
Comorbidities					
Diabetes		196 (43.5)	164 (42.5)	32 (49.2)	0.310
Hypertension		169 (37.5)	139 (36.0)	30 (46.2)	0.118
Chronic obstructive pulmonary disease		15 (3.3)	10 (2.6)	5 (7.7)	0.050
Obesity ≥ 30		31 (6.9)	25 (6.5)	6 (9.2)	0.426
Chronic kidney disease		17 (3.8)	11 (2.9)	6 (9.2)	0.024
Coronary artery disease		31 (6.9)	23 (6.0)	8 (12.3)	0.106
Cerebrovascular disease		4 (0.9)	3 (0.8)	1 (1.5)	0.465
Symptoms at admission					
Shortness of breath		413 (91.6)	351 (90.9)	62 (95.4)	0.232
Fever		158 (35.0)	128 (33.2)	30 (46.2)	0.042
Cough		309 (68.5)	259 (67.1)	50 (76.9)	0.115
Fatigue		354 (78.7)	301 (78.2)	53 (81.5)	0.541
Severity of illness score					
SOFA score at admission*		2.40 ± 1.06	2.30±0.93	3.05±1.49	<0.001
Treatment					
Vasopressor		18 (4.0)	1 (0.3)	17 (26.6)	<0.001
Non-Invasive Ventilation (NIV)		446 (98.9)	383 (99.2)	63 (96.9)	0.101
Invasive ventilation		38 (8.4)	4 (1.04)	34 (52.31)	<0.001
Interval between symptoms onset to admission ‡		4 (3, 7)	4 (3, 7)	4 (3, 6)	0.996
Duration of respiratory support days ‡		6 (4, 10)	6 (4, 9.5)	6 (3, 10)	0.688
Duration of invasive ventilation days ‡		1 (1, 3)	12 (2, 14)	1 (1, 3)	0.019
Duration of hospital stay days ‡		14 (10, 18)	14 (11, 19)	8 (5, 14)	<0.001

‡ Median (IQR) days in days – Mann Whitney U test was used

*Mean ± SD – Independent t test was used

PLACID TRIAL Mortality Assessment

Table 2: Univariate and Multivariable Cox-regression for baseline characteristics, Laboratory parameters and Inflammatory biomarkers

Variables		Univariate Cox regression			Multivariable Cox regression (Model A)			Multivariable Cox regression (Model B)		
		HR	95% CI	P value	HR	95% CI	P value	HR	95% CI	P value
Age	≤40	1.00			1.00			1.00		
	41-59	2.04	0.90 – 4.65	0.089	1.25	0.51 – 3.09	0.623	1.55	0.66 – 3.63	0.310
	≥60	3.51	1.53 – 8.06	0.003	1.42	0.57 – 3.54	0.456	1.72	0.71 – 4.19	0.231
Gender	Male	1.19	0.65 – 2.18	0.581						
Blood Group	O	1.00								
	A	0.94	0.47 – 1.88	0.868						
	B	1.13	0.63 – 2.02	0.693						
	AB	2.01	0.81 – 4.97	0.132						
Comorbidities	No Comorbidities	1.00			1.00			1.00		
	1	1.62	0.81 – 3.24	0.172	1.22	0.60 – 2.49	0.586	1.31	0.60 – 2.85	0.503
	2 or More	2.25	1.17 – 4.32	0.014	1.78	0.90 – 3.51	0.095	2.66	1.29 – 5.51	0.008
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00					
	5-10	4.88	1.82 – 13.06	0.002	3.24	1.19 – 8.80	0.021			
	>10	27.49	11.67 – 64.75	<0.001	9.38	3.67 – 23.99	<0.001			
Platelet count ‡ (* 10 ⁹ /L)	<100	6.53	3.10 – 13.75	<0.001						
	≥ 100	1.00								
SOFA score †		1.61	1.48 – 1.75	<0.001	1.21	1.07 – 1.36	0.002			
D-dimer(mg/L) §	<0.5	1.00						1.00		
	0.5 - 1.0	1.53	0.63 – 3.68	0.347				1.29	0.53 – 3.14	0.568
	>1.0	3.35	1.55 – 7.23	0.002				2.51	1.14 – 5.51	0.022
Ferritin(ng/mL) §	<500	1.00						1.00		
	≥500	4.09	2.31 – 7.23	<0.001				2.66	1.46 – 4.85	0.001
CRP§ (mg/L)		1.0003	0.999 – 1.001	0.150						
LDH§ (IU/L)	<450	1.00						1.00		
	≥ 450	4.88	2.73 – 8.72	<0.001				2.96	1.61 – 5.45	0.001
PaO2/FiO2†	<100 (severe)	23.11	12.81 – 41.69	<0.001	3.37	1.54 – 7.41	0.002			
	100-200(moderate)	5.63	2.97 – 10.68	<0.001	1.84	0.88 – 3.82	0.103			
	>200 (Mild)	1.00			1.00					
Interval from onset of symptoms to admission		1.05	0.97 – 1.12	0.228						
Vasopressor support		11.36	6.47 – 19.96	<0.001						
Invasive ventilation support		19.57	11.81 – 32.41	<0.001						

†Laboratory Parameters were measured at day 0,1,3,5,7 and day 14: § Inflammatory biomarkers values were measured at day 0,3 and day 7
(Model A) Multivariable Cox model for Age, comorbidities with Laboratory Parameters, (Model B) Multivariable Cox model for Age, comorbidities with inflammatory biomarker values

PLACID TRIAL Mortality Assessment

447

For peer review only

PLACID TRIAL Mortality Assessment

Figure 1:





Figure 1: Line graph for SOFA score showing difference between survivors and non survivors

PLACID TRIAL Mortality Assessment**Supplementary****Flowchart for the study protocol**

BMJ Open

Factors associated with mortality among moderate and severe COVID 19 patients – secondary analysis of a Randomized Controlled Trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-050571.R1
Article Type:	Original research
Date Submitted by the Author:	16-Jun-2021
Complete List of Authors:	<p>Mammen, Joy; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Kumar, Snehil; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Thomas, Lovely; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Kumar, Gunjan ; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Zachariah, Anand; Christian Medical College and Hospital Vellore, Medicine</p> <p>Jeyaseelan, Lakshmanan; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Peter, John Victor; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Agarwal, Anup; ICMR, Clinical Trial and Health Systems Research Unit</p> <p>Mukherjee, Aparna; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Chatterjee, Pranab; ICMR, Translational Global Health Policy and Research Cell</p> <p>Bhatnagar, Tarun ; National Institute of Epidemiology, ICMR School of Public Health</p> <p>Rasalam, Jess; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Chacko, Binila; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Mani, Thenmozhi; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Joy, Melvin; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Rupali, Priscilla; Christian Medical College and Hospital Vellore, Infectious Diseases</p> <p>Murugesan, Malathi; Christian Medical College and Hospital Vellore, Hospital Infection Control Committee</p> <p>Daniel, Dolly; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Latha, B; Madras Medical College, Transfusion Medicine</p> <p>Bundas, Sunita; SMS Medical College and Hospital, Transfusion Medicine</p> <p>Kumar, Vivek; Sir HN Reliance Foundation Hospital and Research Centre, Critical Care</p> <p>Dosi, Ravi; Sri Aurobindo Institute of Medical Sciences, Respiratory Medicine</p>

	Khambholja, Janakkumar; Smt NHL Municipal Medical College, Internal Medicine de Souza, Rosemarie; BYL Nair Charitable Hospital, Department of Medicine Chander, B; Gandhi Medical College and Hospital, Internal Medicine Bahadur, Shalini; Government Institute of Medical Sciences, Pathology Dube, Simmi; Gandhi Medical College Bhopal, Internal Medicine Suri, Amit; Atal Bihari Vajpayee Institute of Medical Sciences and Ram Manohar Lohia Hospital, Pulmonary Medicine Jindal, Aikaj ; Satguru Pratap Singh Hospitals, Transfusion Medicine Shrivastav, Om ; Kasturba Hospital for Infectious Diseases, Infectious diseases Barge, Vijay; RSCM Government Medical College, Medicine Bajpayee, Archana; AIIMS Jodhpur, Transfusion Medicine Malhotra, Pankaj; Post Graduate Institute of Medical Education and Research, Department of Internal Medicine Singh, Neha; AIIMS Patna, Transfusion Medicine Tambe, Muralidhar; Department of Community Medicine, B J Govt. Medical College Sharma, Nimisha; ESIC Medical College and Hospital Faridabad, Transfusion Medicine Bhat, Shreepad; Smt Kashibai Navale Medical College and General Hospital, Internal Medicine Kaulgud, Ram; Karnataka Institute of Medical Sciences, Internal Medicine Gurtoo, Anil; Lady Hardinge Medical College, Internal Medicine Reddy, Himanshu; King George Medical College, Internal Medicine Upadhyay, Kamlesh; Byramjee Jeejabhoy Medical College, Internal medicine Jain, Ashish ; Mahatma Gandhi Medical College and Hospital, Respiratory Medicine Patel, Tinkal; Government Medical College, Internal Medicine Nagori, Irfan; GMERS Medical College Gotri Vadodara, Medicine Jha, Pramod ; Sumandeep Vidyapeeth University, Internal Medicine Babu, KV; Sri Venkateswara Institute of Medical Sciences, Transfusion Medicine Aparna, C; Kurnool Medical College Panjwani, Sunil; Government Medical College, Internal Medicine Natarajan, M; Madurai Medical College, Internal Medicine Baldi, Milind; Mahatama Gandhi Memorial Medical College, Internal Medicine Khadke, Vrushali; Poona Hospital and Research Centre, Interventional Pulmonology Dua, Seema; Super Speciality Paediatric Hospital and Teaching Hospital, Transfusion Medicine Singh, Ravindraa; Aditya Birla Memorial Hospital, Transfusion Medicine Sharma, Ashish; R. D. Gardi Medical College, Medicine Sharma, Jayashree; Seth GS Medical College and KEM Hospital, Transfusion Medicine Gokhale, Yojana; Lokmanya Tilak Municipal Medical College and General Hospital, Internal Medicine Yadav, Pragya; National Institute of Virology, Maximum Containment Laboratory, ICMR Sapkal, Gajanan; National Institute of Virology, Diagnostic Virology Group, ICMR Kaushal, Himanshu; National Institute of Virology, Human Influenza Group, ICMR Kumar, Saravana; National Institute of Epidemiology, Epidemiology and Biostatistics Division
Primary Subject	Public health

Heading:	
Secondary Subject Heading:	Public health
Keywords:	COVID-19, Public health < INFECTIOUS DISEASES, Adult intensive & critical care < INTENSIVE & CRITICAL CARE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

PLACID TRIAL Mortality Assessment

1 Factors associated with mortality among moderate and severe COVID 19 patients – 2 secondary analysis of a Randomized Controlled Trial

3 Joy J Mammen, Snehil Kumar, Lovely Thomas, Gunjan Kumar, Anand Zachariah, Lakshmanan
4 Jeyaseelan, John Victor Peter, Anup Agarwal, Aparna Mukherjee, Pranab Chatterjee, Tarun
5 Bhatnagar, Jess Elizabeth Rasalam, Binila Chacko, Thenmozhi Mani, Melvin Joy, Priscilla Rupali,
6 Malathi Murugesan, Dolly Daniel, B Latha , Sunita Bundas, Vivek Kumar, Ravi Dosi, Janakkumar R
7 Khambholja, Rosemarie de Souza, B Thrilok Chander, Shalini Bahadur, Simmi Dube, Amit Suri,
8 Aikaj Jindal, Om Shrivastav, Vijay Barge, Archana Bajpayee, Pankaj Malhotra, Neha Singh,
9 Muralidhar Tambe, Nimisha Sharma, Shreepad Bhat, Ram S Kaulgud, Anil Gurtoo, D Himanshu
10 Reddy, Kamlesh Upadhyay, Ashish Jain, Tinkal C Patel, Irfan Nagori, Pramod R Jha, KV Sreedhar
11 Babu, C Aparna, Sunil Jodharam Panjwani, M Natarajan, Milind Baldi, Vrushali Khirid Khadke,
12 Seema Dua, Ravindraa Singh, Ashish Sharma, Jayashree Sharma, Yojana A Gokhale, Pragya D
13 Yadav, Gajanan Sapkal, Himanshu Kaushal, V Saravana Kumar

17 Corresponding Author:

18 Joy J Mammen MD

19 Professor & Head

21 Address

22 Department of Transfusion Medicine and Immunohaematology

23 Christian Medical College Vellore

24 Tamil Nadu, India

25 Pin code 632004

27 E mail joymammen@cmcvellore.ac.in

29 Phone +91 416 228 2536

31 Fax +91 416 223 2035

PLACID TRIAL Mortality Assessment

32

Name	Designation and affiliation	Full Address
Joy J Mammen	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Snehil Kumar	Senior Resident, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lovely Thomas	Assistant Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Gunjan Kumar	Scientist C, Clinical Trial & Health Systems Research Unit, Indian Council of Medical Research (ICMR)	New Delhi, India.
Anand Zachariah	Professor, Department of Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lakshmanan Jeyaseelan	Professor, Clinical Epidemiology Unit, Christian Medical College	Vellore, Tamil Nadu, India.
John Victor Peter	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Anup Agarwal	Consultant, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Aparna Mukherjee	Scientist E, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Pranab Chatterjee	Scientist B, Translational Global Health Policy and Research Cell, ICMR	New Delhi, India.
Tarun Bhatnagar	Scientist E, ICMR School of Public Health, ICMR -National Institute of Epidemiology	Chennai, Tamil, India
Jess Elizabeth Rasalam	Fellow, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Binila Chacko	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Thenmozhi Mani	Senior Demonstrator, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Melvin Joy	Associate Research Officer, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Priscilla Rupali	Professor, Department of Infectious Diseases, Christian Medical College	Vellore, Tamil Nadu, India.
Malathi Murugesan	Assistant Professor, Hospital Infection Control Committee, Christian Medical College	Vellore, Tamil Nadu, India.
Dolly Daniel	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
B Latha	Professor, Department of Transfusion Medicine, Madras Medical College	Chennai, Tamil Nadu, India
Sunita Bundas	Professor, Department of Immunohaematology and Transfusion Medicine, SMS Medical College and Hospital	Jaipur, Rajasthan, India
Vivek Kumar	Professor, Department of Critical Care Medicine, Sir H N Reliance Foundation Hospital	Mumbai, Maharashtra, India
Ravi Dosi	Professor, Department of Respiratory Medicine, Sri Aurobindo Institute of Medical Sciences	Indore, Madhya Pradesh, India
Janakkumar R Khambholja	Professor, Department of Internal Medicine, Smt NHL Municipal Medical College	Ahmedabad, Gujarat, India
Rosemarie de Souza	Professor, Department of Internal Medicine, BYL Nair Charitable Hospital & TN Medical College	Mumbai, Maharashtra, India
B Thrilok Chander	Professor, Department of Internal Medicine, Gandhi Medical College, Secunderabad	Secunderabad, Telangana, India
Shalini Bahadur	Associate Professor, Department of Pathology, Government Institute of Medical Sciences	Greater Noida, Uttar Pradesh, India
Simmi Dube	Professor, Department of Internal Medicine, Gandhi Medical College	Bhopal, Madhya Pradesh, India
Amit Suri	Professor, Department of Pulmonary Medicine, Atal Bihari Vajpayee Institute of Medical Sciences and RML Hospital	New Delhi, India
Aikaj Jindal	Professor, Department of Transfusion Medicine, Satguru Partap Singh Hospitals	Ludhiana, Punjab, India
Om Shrivastav	Professor, Department of Infectious Diseases, Kasturba Hospital	Mumbai, Maharashtra, India
Vijay Barge	Professor, Department of Medicine, RSCM Government Medical College	Kolhapur, Maharashtra, India
Archana Bajpayee	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Jodhpur, Rajasthan, India
Pankaj Malhotra	Professor, Department of Internal Medicine, Post Graduate Institute of Medical Education and Research	Chandigarh, India
Neha Singh	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Patna, Bihar, India
Muralidhar Tambe	Professor, Department of Community Medicine Byramjee Jeejeebhoy Medical College	Pune, Maharashtra, India

PLACID TRIAL Mortality Assessment

Nimisha Sharma	Professor, Department of Immunohematology and blood transfusion, ESIC Medical College and Hospital	Faridabad, Haryana, India
Shreepad Bhat	Professor, Department of Internal Medicine, Smt. Kashi Bai Navale Medical College	Pune, Maharashtra, India
Ram S Kaulgud	Professor, Department of Internal Medicine, Karnataka Institute of Medical Sciences	Hubali, Karnataka, India
Anil Gurtoo	Professor, Department of Internal Medicine, Lady Hardinge Medical College	New Delhi, India
D Himanshu Reddy	Professor, Department of Internal Medicine, King George's Medical University	Lucknow, Uttar Pradesh, India
Kamlesh Upadhyay	Professor, Department of Internal Medicine, Byramjee Jeejeebhoy Medical College	Ahmedabad, Gujarat, India
Ashish Jain	Professor, Department of Respiratory Medicine, Mahatma Gandhi University of Medical Sciences and Technology	Jaipur, Rajasthan, India
Tinkal C Patel	Professor Department of Internal Medicine, Government Medical College	Surat, Gujarat, India
Irfan Nagori	Professor, Department of Medicine, GMERS Medical College and Hospital	Vadodara, Gujarat, India
Pramod R Jha	Professor, Department of Internal Medicine, Sumandeep, Vidyapeeth and Dhiraj Hospital	Vadodara, Gujarat, India
KV Sreedhar Babu	Professor, Department of Transfusion Medicine, Sri Venkateswra Institute of Medical Sciences	Tirupati, Andhra Pradesh, India
C Aparna	Professor, Department of Pathology, Kurnool Medical College	Kurnool, Andhra Pradesh, India
Sunil Jodharam Panjwani	Professor, Department of Internal Medicine, Government Medical College	Bhavnagar, Gujarat, India
M Natarajan	Professor, Department of Internal Medicine, Madurai Medical College	Madurai, Tamil Nadu, India
Milind Baldi	Professor, Department of Respiratory Medicine, Mahatma Gandhi Memorial Medical College	Indore, Madhya Pradesh, India
Vrushali Khirid Khadke	Professor, Department of Interventional Pulmonology, Poona Hospital and Research Center	Pune, Maharashtra, India
Seema Dua	Professor, Department of Transfusion Medicine, Super Specialty Paediatric Hospital and Post Graduate Teaching Institute	Noida, Uttar Pradesh, India
Ravindraa Singh	Professor, Department of Transfusion Medicine, Aditya Birla Memorial Hospital	Pune, Maharashtra, India
Ashish Sharma	Professor, Department of Medicine, RD Gardi Medical College	Ujjain, Madhya Pradesh, India
Jayashree Sharma	Professor, Department Transfusion Medicine, Seth GS Medical College and KEM Hospital	Mumbai, Maharashtra, India
Yojana A Gokhale	Professor, Department of Internal Medicine, Lokmanya Tilak Municipal Medical College & General Hospital	Mumbai, Maharashtra, India
Pragya D Yadav	Scientist E, Maximum Containment Laboratory, ICMR – National Institute Virology	Pune, Maharashtra, India
Gajanan Sapkal	Scientist E, Diagnostic Virology Group, ICMR – National Institute Virology	Pune, Maharashtra, India
Himanshu Kaushal	Scientist B, Human Influenza group, ICMR – National Institute Virology	Pune, Maharashtra, India
V Saravana Kumar	Scientist B, Epidemiology and Biostatistics Division, ICMR – National Institute of Epidemiology	Chennai, Tamil Nadu, India

33

PLACID TRIAL Mortality Assessment

Abstract

Objective:

Large data on the clinical characteristics and outcome of COVID-19 in the Indian population is scarce. We analyzed the factors associated with mortality in a cohort of moderate to severely ill COVID-19 patients enrolled in a randomized trial on convalescent plasma.

Setting:

39 public and private hospitals across India.

Participants:

Of the 464 patients recruited, two were lost to follow-up, nine withdrew consent and two patients did not receive the intervention after randomization. The cohort of 451 participants with known outcome at 28-days was analyzed.

Design:

Secondary analysis of data from a Phase II, Open Label, Randomized Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID TRIAL).

Primary outcome measure:

Factors associated with all-cause mortality at 28-days post-enrolment.

Results:

The mean (SD) age was 51±12.4 years; 76.7% were males. Admission SOFA score was 2.4±1.1. Non-invasive ventilation, invasive ventilation and vasopressor therapy were required in 98.9%, 8.4% and 4.0% respectively. The 28-day mortality was 14.4%. Median time from symptom onset to hospital admission was similar in survivors (4 days; IQR 3-7) and non survivors (4 days; IQR 3-6). Patients with two or more co-morbidities had 2.25 (95%CI: 1.18-4.29, p=0.014) times risk of death. When compared with survivors, admission IL-6 levels were higher (p<0.001) in non-survivors and increased further on Day 3. On multivariable Fine and Gray model , severity of illness (SHR 1.22, 95%CI:1.11-1.35,p<0.001), PaO₂/FiO₂ ratio <100 (3.47, 1.64-7.37, p=0.001), Neutrophil Lymphocyte ratio (NLR) >10 (9.97, 3.65-27.13, p<0.001), D-dimer >1.0mg/L (2.50,1.14-5.48, p=0.022), ferritin ≥500ng/mL (2.67, 1.44-4.96, p=0.002) and LDH ≥450 IU/L (2.96, 1.60-5.45, p=0.001) were significantly associated with death.

Conclusion:

In this cohort of moderate to severely ill COVID-19 patients, severity of illness, underlying co-morbidities and higher levels of inflammatory markers were significantly associated with death.

PLACID TRIAL Mortality Assessment

Trial Registration:

The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775).

Article Summary

Strengths and limitations of this study

Strengths

There is no study from India, with representation from multiple states that has detailed the clinical profile and evaluated for factors associated with death. This study may help with strategic planning at a national level.

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the present analysis need not be adjusted for convalescent plasma intervention.

There may be variability of treatment provided in the multiple centres, however, care was taken that patients received best standard of care for COVID-19 dictated by the best available evidence at the time and guidelines for the management of COVID-19 issued by health authorities of the Indian government.

Limitations

The laboratory and biomarker assays for ferritin, lactate dehydrogenase, C reactive protein, and D-dimer were conducted using tests from different manufacturers.

Participants of this study may not comprise a true observational cohort as this was a post hoc analysis of a randomized control trial data, also our study did not analyse the effect of SARS-CoV-2 variants causing a high mortality in younger population during the second wave of COVID-19 infection, therefore extrapolation to the general population must be carefully qualified.

PLACID TRIAL Mortality Assessment

93 Introduction

94 The first human case of Corona Virus Disease 19 (COVID-19) caused by the novel coronavirus
95 (named Severe Acute Respiratory Syndrome Coronavirus 2, SARS-CoV-2) was reported in Wuhan
96 City, China in December 2019. On 30 January 2020, the World Health Organization (WHO)
97 declared that the outbreak of COVID-19 constituted a Public Health Emergency of International
98 Concern (1). Based on the high levels of global spread and the severity of COVID-19, on 11 March
99 2020, the Director-General of the WHO declared the COVID-19 outbreak a pandemic (2). The
100 sudden outbreak followed by rapid spread in a globalized world, resulted in a major medical
101 burden, besides affecting socio-economic well-being among all nations.

102 In India, the disease was first detected on 30 January 2020 in the state of Kerala, in a student who
103 returned from Wuhan (3). After a brief, initial respite, the virus has spread at a rapid pace in India,
104 resulting in more than 10 million confirmed cases as of December, 2020 with more than 145,000
105 deaths (4).

106 Patients diagnosed with COVID-19 have primarily respiratory symptoms. Most patients diagnosed
107 with COVID-19 experience mild to moderate respiratory illness, fever, dry cough, fatigue and
108 recover without requiring special treatment (5). Oxygen desaturation is the hallmark of
109 progression. Patients with underlying medical problems like cardiovascular disease, diabetes,
110 chronic respiratory disease, and cancer are more likely to develop serious illness. These patients
111 may develop viral pneumonia, with resultant dyspnea and hypoxemia which may progress to
112 respiratory or multi- system failure and even death (6). There is paucity of large-scale data on the
113 clinical characteristics, outcomes of COVID-19 in the Indian population and evaluation of risk
114 factors with an unfavorable outcome at a national level. Identification of such potential risk factors
115 is important to anticipate medical treatment and to reduce the mortality burden for severe COVID-
116 19 illness by proactive interventions.

117 The Indian Council of Medical Research (ICMR) conducted a randomized trial (A Phase II, Open
118 Label, Randomized Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to
119 Limit COVID-19 Associated Complications in Moderate Disease, PLACID TRIAL) to determine the
120 effectiveness and safety of convalescent plasma in patients with moderate to severely ill COVID-19
121 to limit progression of disease (7). Patients received standard of care for COVID-19 in keeping with
122 the institutional protocols, based on the best available evidence at the time and guidelines for the
123 management of COVID-19 issued by national health authorities. Participants in the intervention

PLACID TRIAL Mortality Assessment

arm received two doses of 200 mL of convalescent plasma, transfused 24 hours apart, in addition to standard of care. The control arm did not receive any additional therapy. The study concluded that the use of convalescent plasma was not associated with a reduction in 28-day mortality (7).

The aim of this analysis was to identify risk factors associated with mortality by mining the data collected from the cohort enrolled in the PLACID TRIAL (7).

Methods

Participants

The study enrolled patients from 39 different hospitals, of which, 29 were teaching public hospitals and 10 were private facilities spread across 14 states and union territories. Patients over the age of 18 years who were confirmed to have COVID-19 based on a positive SARS-CoV-2 RT-PCR test and moderate to severely ill with either a partial pressure of oxygen in arterial blood/fraction of inspired oxygen ($\text{PaO}_2/\text{FiO}_2$) ratio between 200-300 or respiratory rate $>24/\text{min}$ and decreased oxygen saturation on room air ($\text{SpO}_2 < 93\%$) were included during the study period from 22 April to 14 July 2020. Patients were followed up for 28-days and assessed for their health status and all-cause mortality. Ethics approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals. Written consent was obtained from patients or their families before enrolling in the study.

Data

Data was obtained from the ICMR PLACID TRIAL database collected in structured paper case record forms and entered in Research Electronic Data Capture system (REDCap, version 8.5 Vanderbilt University, TN). The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775). After trial completion, based on cooperative agreement between the centers, and IRB permission, the data was shared and analyzed further, to explore for other meaningful results. No separate ethical clearance was taken for this study.

Demographic, clinical, laboratory tests and outcome data were collected prospectively. Clinical symptoms need for organ support (respiratory, hemodynamic), laboratory tests (complete blood count, coagulation profile, serum biochemical profile, renal and liver function tests) were monitored serially on day of enrollment (day 0) and on day 1, 3, 5, 7, 14 and 28. Inflammatory

PLACID TRIAL Mortality Assessment

biomarkers [lactate dehydrogenase (LDH), serum ferritin, and C-reactive protein (CRP)] were tested at admission and on day 3 and 7 whereas; Interleukin 6 (IL-6) was done at admission and on day 3.

The outcome of interest was all-cause day 28 mortality. We evaluated for association between laboratory parameters and mortality.

Statistical Methods:

Mean and standard deviation (SD) or median and inter-quartile range (IQR) were used for continuous variables as appropriate, and categorical variables number and proportions were used. To find the association between mortality and study variables, Chi-square test/Fisher's exact test were used. To find the mean difference across the groups, independent t test was used. Similarly, Mann Whitney U test was used to compare median difference. The end point of interest was all-cause mortality (event of interest) at day 28 from the time of enrollment, discharged alive (competing event) and hospital admission after day 28 (censored), whichever is earlier. Discharged alive was treated as a competing event because the event of discharged alive precludes the event of all-cause mortality. Clinically important baseline variables and time dependent covariates were included in multivariable Fine and Gray regression model for competing endpoints and sub-distribution hazard ratios were presented. Two multivariable models were developed. The first model included clinical and laboratory parameters tested on day 0, 1, 3, 5, 7, 14 and 28 while the second included inflammatory biomarkers tested on day 0, 3 and 7 after adjusting for age and comorbidities. For certain laboratory markers such as D-dimer, ferritin and LDH, clinically relevant thresholds were used for the analysis rather than using these data as continuous variables. The clinically relevant thresholds for these variables were set as >1.0 mg/L for D-dimer, ≥500 mg/mL for Ferritin and ≥450 IU/L for LDH. Variables included parameters that were strongly associated with mortality at univariate analysis or known from previous literature to be strongly associated with outcome. The model assumption was verified using log-log S (t) plots and Global test. A p-value of less than 0.05 levels was considered as statistically significant. All statistical analyses were performed using STATA version 16.0 (StataCorp. 2019. College Station, TX).

PLACID TRIAL Mortality Assessment

Patient and public involvement

Patients and the public were not involved in the design, conduct, reporting, or dissemination plans of our research. The study results will be disseminated to the study participants via their treating doctors.

Results

The PLACID Trial recruited 464 eligible patients for the study. The primary outcome at 28-days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomization and two patients did not receive the intervention after randomization as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients (*supplementary*).

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the analysis did not adjust for convalescent plasma intervention. The distribution of patients in intervention and control arms were 50.3% (n=227) and 49.7% (n=224), respectively. The mean (SD) age of the cohort was 51 ± 12.4 years; 76.7% were males. Table 1 shows distribution of demographic variables and clinical parameters in the study population.

The most common presenting symptoms were shortness of breath (91.6%), fatigue (78.7%), cough (68.5%) and fever (35%). Comorbidities were present in 59.9% of patients; 31.7% had any one comorbidities and 28.2% had two or more comorbidities. The most frequent comorbidities were diabetes (43.5%), hypertension (37.5%), obesity (6.9%) and Chronic Obstructive Pulmonary Disease (COPD) (3.3%). There was history of smoking in 8.2%. The time from onset of symptoms to admission was four days (IQR 3-7 days). Majority of the patients required non-invasive (98.9%) ventilatory support. The median duration of respiratory support was six days (IQR 4-10 days). In this cohort, 4% patients required vasopressor support. None of the patients required Extra Corporeal Membrane Oxygenation (ECMO) or dialysis support.

The all-cause mortality at 28-days was 14.4% (95%CI: 11.5-17.9, n=65). Median time from symptom onset to hospital admission was four days in survivors (IQR 3-7 days) and non survivors (IQR 3-6 days). The frequency of shortness of breath, cough and fatigue were similar in survivors and non survivors; however, the presence of fever at admission was significantly (p=0.042) associated with death (table 1). Other than COPD and CKD (chronic kidney disease), other

PLACID TRIAL Mortality Assessment

comorbidities were not significantly associated with death (table 1). Admission Sequential Organ Failure Assessment (SOFA) score was higher in non survivors. The need for invasive mechanical ventilation, duration of invasive mechanical ventilation and vasopressor therapy were associated with death (table 1).

On univariate analysis (table 2), there was an association between increasing age and mortality. Patients with two or more comorbidities had a 2.25 (95%CI: 1.18-4.29, p=0.014) times increased chance of mortality. There was a strong mortality association for platelet count $<100 \times 10^9/L$ (SHR 6.88, 95%CI: 3.61-13.13, p<0.001), neutrophil lymphocyte ratio (NLR) >10 (28.84, 11.92-69.76, p<0.001), LDH ≥ 450 IU/L (4.88, 2.72-8.75, p<0.001), D-dimer >1 mg/L (3.34, 1.55-7.19, p=0.002) and ferritin ≥ 500 ng/mL (4.11, 2.28-7.41, p<0.001). Admission IL-6 levels were significantly (p<0.001) higher (76.00, 18.27-171.77) in non survivors than in survivors (18.51, 4.26-56.86). By day 3, IL-6 levels dropped to 11.6 (2.64-45.84) in survivors while it nearly doubled in non-survivors (140.35, 21.56-427.36). Univariate analysis of CRP did not show any statistical significance (1.0003, 0.999-1.001, p=0.080).

The need for invasive ventilation and vasopressors were associated with death (table 2). Increasing SOFA score was associated with mortality (1.63, 1.54-1.74, p<0.001). The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score progressively increased between the two groups over time (figure 1). Mortality proportionately also increased with lower PaO₂/FiO₂ values with sub-distribution hazard ratio of 25.64 (14.8-44.41, p<0.001) in the severe group as compared to the mild group.

Two models were run for multivariable Fine and Gray regression model over a period of time (table 3). Model A included age, comorbidities, PaO₂/FiO₂, NLR and SOFA score. Model A revealed significant sub-distribution hazard ratios for PaO₂/FiO₂ ratio <100 (3.47, 1.64-7.37, p=0.001), NLR >10 (9.97, 3.65-27.13, p<0.001), SOFA score (1.22, 1.11- 1.35, p<0.001) after adjusting for age and comorbidities. Model B included age, comorbidities, D-dimer, ferritin and LDH. D-dimer >1 mg/L (2.50, 1.14-5.48, p=0.022), ferritin ≥ 500 ng/mL (2.67, 1.44-4.96, p=0.002) and LDH ≥ 450 IU/L (2.96, 1.60-5.45, p=0.001), were associated with mortality after adjusting for age and comorbidities (table 3). IL-6 was omitted from the model as it was not measured on Day 7.

PLACID TRIAL Mortality Assessment**Discussion**

In this study that enrolled patients from across India, we were able to identify clinical, biomarkers (D-dimer, LDH, ferritin) and SOFA score as factors that could indicate increased risk of death in moderately to severely ill COVID-19 patients from PLACID trial cohort. The definition of clinical grading of severity is different in India as compared to other countries (8–12). Mortality of critically ill COVID-19 patients varies significantly among the published case series, ranging from 16% to 78% (13–19). Similarly two studies from Wuhan, which included moderately as well as critically ill patients, have shown mortality rates of 3.77% and 14.14% (20,21). This wide variability can be explained by differences in the age of the population, distribution of risk factors, health system response across different countries, varied treatment protocols and follow-up. In a series of critically ill patients in China, 28-day ICU mortality was 61.5% (22). In a multicentric study from Italy, the mortality risk for patients without respiratory failure at admission was of 1% after 15 days while survival in patients with moderate-to-severe respiratory failure ($\text{PaO}_2/\text{FiO}_2 \leq 200$ mm Hg) at admission was only 56% at 15 days (23). The fatality rate reported in Europe and the United States of America is significantly higher than in China (24). Therefore, findings obtained in a specific country might not be automatically extrapolated and national cohorts must be studied.

In our study population, mortality increased with age, which is similar to the pattern observed in other countries affected by COVID-19. Age seems to affect the time from hospitalization to death. Age-specific death rates was quite similar in studies from Asia, Europe and North America (25). South Korea, Italy, France, Germany, England and Wales, and Spain COVID-19 attributed mortality rates rise by about 12%/year while the United States and Wuhan, China showed a slower rate of increase about 9.5%/year of age (26). In a meta-analysis of 61, 11,583 subjects, 23.2% of patients were aged ≥ 80 years and showed an average mortality rate of 12.10%, the lowest being in China (3.1%) and the highest in the United Kingdom (20.8%) and New York State (20.99%). In the same study, highest mortality rate was observed in patients aged ≥ 80 years. The largest increase in mortality risk was observed in patients aged 60 to 69 years compared with those aged 50 to 59 years (odds ratio 3.13, 95% CI: 2.61–3.76) (27).

Presence of comorbidities significantly increases the death risk of COVID-19. A higher risk of mortality was seen in our patients who had CKD and COPD. Meta-analysis, including 1389 COVID-19 patients with 19.7% having severe disease showed a significant association of CKD with severe COVID-19 with pooled odds ratio as 3.03 (28). Similarly, the estimated mortality risk in patients

PLACID TRIAL Mortality Assessment

with COPD was three times of those without ($p<0.05$) (29). We found that 43.5% of our patients had diabetes which is markedly higher as compared to patients from Korea which showed that 16.97% had diabetes mellitus (30). Our analysis showed that the presence of diabetes was not significantly different between survivors and non survivors (42.5% vs. 49.2%, $p=0.310$), in contrast to the study from South Korea (30) which showed a much higher mortality among diabetic patients than in those without (20.0% vs. 4.8%). Hypertension and obesity were not significantly different among survivors and non-survivors in our study. However, the presence of two or more comorbidities was associated with mortality in our study.

Univariate Fine-Gray model identified several other prognostic markers for mortality, most notably age ≥ 60 years, PaO_2/FiO_2 ratio <100 , NLR >10 , platelet count $<100 \times 10^9/L$, ferritin $>500ng/mL$, LDH $>450 IU/L$ and D-dimer $>1mg/L$. Our study showed similar findings when compared with studies from Wuhan (31). Older age, leukocytosis, and high LDH level have been reported to be risk factors associated with in-hospital death in other studies also (32–34). IL-6 levels were significantly different in survivors and non-survivors at admission. By day 3 survivors had reducing IL-6 while it nearly doubled in non survivors.

Mortality was higher among patients requiring invasive mechanical ventilation (SHR 19.57, 12.21–31.35, $p<0.001$) and those requiring vasopressors (SHR 11.36, 7.79–16.56, $p<0.001$). However, the median duration of invasive ventilation for survivors was 12 days (2, 14) and that for non-survivors was one day (1, 3). These results suggest that patients with acute respiratory failure from COVID-19 may recover, even with severe disease requiring longer ventilation, and that probably the sickest patients die early reflecting lower duration of invasive ventilation in non survivors. Therefore, invasive ventilation should be timely and effectively provided.

In our study, the SOFA score was recognized as a valuable tool that could be used to prognosticate outcome of patients with COVID-19. Univariate and multivariable competing risk regression models showed that the increase in SOFA score was related to mortality, with a clearly divergent pattern between the two groups. Thus, an increasing SOFA score over time may be a factor that can be used to identify a subset of patients who may have an unfavorable outcome. Studies have shown that the SOFA score could be used to evaluate severity and 60-day mortality of COVID-19 with the optimal cut-off score of 5 (35).

PLACID TRIAL Mortality Assessment

Limitations of the study include the variability of treatment provided in the multiple centers. The participants of this study may not comprise a true observational cohort as this was a post hoc analysis of a randomized control trial data and extrapolation to the general population must be carefully qualified. Our study did not analyse the effect of SARS-CoV-2 variants causing a high mortality in younger population during the second wave of COVID-19 infection and this may limit generalizability of the data to the second wave. Despite these limitations, this study provides a comprehensive overview of prognostic factors in moderate to severely ill COVID-19 patients that included patients from across the country.

Conclusion

A favorable outcome can be expected in moderate to severely ill COVID-19 patients. Older age, multiple comorbidities, low PaO₂/FiO₂ ratio and deranged inflammatory markers are associated with worse prognosis. Serial SOFA score can be used for prognostication. Understanding symptoms, burden of comorbidities and systematic monitoring of key laboratory parameters offer opportunities for targeted intervention in COVID-19 with the use of anti-inflammatory or immunomodulatory agents.

Figure legend

Figure 1 showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

FOOTNOTES:

Authors Contributions:

Study design: JJM, LJ, JVP

Clinical Management: SK, LT, AZ, JER, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG.

Data collection: JJM, SK, LT, AZ, AA, AM, GK, PC, TB, JER, PR, MM, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG, PDY, GS, HK, VSK.

Data Analysis: JJM, SK, LJ, TM, MJ, PR, MM, DD, JVP.

PLACID TRIAL Mortality Assessment

Data Interpretation: JJM, SK, AZ, LJ, JER, TM, MJ, DD, JVP

Manuscript writing: JJM, SK, LT, AZ, LJ, JER, BC, TM, MJ, PR, MM, DD, JVP, AA, AM, GK, HK, PC, TB

Study Administration: JJM, BC, JVP, AA, AM, GK, HK, PC, TB

Funding support: The funding source for the primary study was ICMR (RFC No. ECD/NTF/20/2020-21/Covid Dated 23.07.2020). However, they had no financial interest in the investigational product. No other author has any competing financial or non-financial interest.

Competing interests: 'None declared'. All authors have completed the ICMJE uniform disclosure form at www.icmje.org.

Ethical Approval: Ethical approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) based in the National Center for Disease Informatics and Research, Indian Council of Medical Research, Bengaluru, Karnataka, as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals.

Data availability statement: Data will be made available, upon request, and must be accompanied by a brief proposal outlining the analysis plan. A signed data access agreement might be needed to ensure data safety and compliance with national rules about data sharing.

Acknowledgement: We acknowledge ICMR for providing us the data collected during PLACID trial which was analyzed in the present study.

References:

1. Mullen L, Potter C, Gostin LO, Cicero A, Nuzzo JB. An analysis of International Health Regulations Emergency Committees and Public Health Emergency of International Concern Designations. *BMJ Glob Health*. 2020 Jun 1;5(6):e002502.
2. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Bio-Medica Atenei Parm*. 2020 19;91(1):157–60.
3. Perappadan BS. India's first coronavirus infection confirmed in Kerala. *The Hindu* [Internet]. 2020 Jan 30 [cited 2020 Nov 16]; Available from: <https://www.thehindu.com/news/national/indias-first-coronavirus-infection-confirmed-in-kerala/article30691004.ece>

PLACID TRIAL Mortality Assessment

4. MoHFW | Home [Internet]. [cited 2021 Jan 3]. Available from: <https://www.mohfw.gov.in/>
5. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol Orlando Fla*. 2020 Jun;215:108427.
6. Wu C, Chen X, Cai Y, Xia J 'an, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med*. 2020 01;180(7):934–43.
7. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre randomised controlled trial (PLACID Trial). *BMJ* [Internet]. 2020 Oct 22 [cited 2020 Nov 16];371. Available from: <https://www.bmj.com/content/371/bmj.m3939>
8. Varghese GM, John R, Manesh A, Karthik R, Abraham OC. Clinical management of COVID-19. *Indian J Med Res*. 2020 May 1;151(5):401.
9. Clinical management of COVID-19 [Internet]. [cited 2021 Jan 3]. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19>
10. Zhang S-Y, Lian J-S, Hu J-H, Zhang X-L, Lu Y-F, Cai H, et al. Clinical characteristics of different subtypes and risk factors for the severity of illness in patients with COVID-19 in Zhejiang, China. *Infect Dis Poverty*. 2020 Jul 8;9(1):85.
11. Xia L, Chen J, Friedemann T, Yang Z, Ling Y, Liu X, et al. The Course of Mild and Moderate COVID-19 Infections—The Unexpected Long-Lasting Challenge. *Open Forum Infect Dis* [Internet]. 2020 Sep 1;7(ofaa286). Available from: <https://doi.org/10.1093/ofid/ofaa286>
12. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis*. 2020 Jun 1;20(6):669–77.
13. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al. Risk Factors Associated With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy. *JAMA Intern Med*. 2020 01;180(10):1345–55.
14. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet Lond Engl*. 2020 15;395(10223):497–506.
15. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, et al. Clinical Course and Outcomes of 344 Intensive Care Patients with COVID-19. *Am J Respir Crit Care Med*. 2020 Apr 8;201(11):1430–4.
16. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA*. 2020 Mar 17;323(11):1061–9.

PLACID TRIAL Mortality Assessment

17. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, et al. Covid-19 in Critically Ill Patients in the Seattle Region - Case Series. *N Engl J Med*. 2020 21;382(21):2012–22.
18. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and Outcomes of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA*. 2020 28;323(16):1612–4.
19. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet Lond Engl*. 2020 28;395(10229):1054–62.
20. Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement, and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect*. 2020 Jun 1;26(6):767–72.
21. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ*. 2020 Mar 26;368:m1091.
22. Yang X, Yu Y, Xu J, Shu H, Xia J 'an, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020 May 1;8(5):475–81.
23. Santus P, Radovanovic D, Sadari L, Marino P, Cogliati C, De Filippis G, et al. Severity of respiratory failure at admission and in-hospital mortality in patients with COVID-19: a prospective observational multicentre study. *BMJ Open [Internet]*. 2020 Oct 10;10(10). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7549463/>
24. Yamamoto N, Bauer G. Apparent difference in fatalities between Central Europe and East Asia due to SARS-COV-2 and COVID-19: Four hypotheses for possible explanation. *Med Hypotheses*. 2020 Nov;144:110160.
25. Demographics of COVID-19 Deaths [Internet]. Ined - Institut national d'études démographiques. [cited 2020 Nov 16]. Available from: <https://dc-covid.site.ined.fr/en/>
26. Goldstein JR, Lee RD. Demographic perspectives on the mortality of COVID-19 and other epidemics. *Proc Natl Acad Sci U S A*. 2020 Sep 8;117(36):22035–41.
27. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583 Subjects. *J Am Med Dir Assoc*. 2020 Jul 1;21(7):915–8.
28. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *Int Urol Nephrol*. 2020 Mar 28;1–2.

PLACID TRIAL Mortality Assessment

29. Venkata VS, Kiernan G. COVID-19 AND COPD: POOLED ANALYSIS OF OBSERVATIONAL STUDIES. *CHEST*. 2020 Oct 1;158(4):A2469.
30. D A, K L, Ds L, Ys L, Ss M. Mortality Rate and Predictors of Mortality in Hospitalized COVID-19 Patients with Diabetes. *Healthc Basel Switz* [Internet]. 2020 Sep 13 [cited 2020 Nov 16];8(3). Available from: <https://europepmc.org/article/med/32933191>
31. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* [Internet]. 2020 May 7;55(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144257/>
32. Li M, Cheng B, Zeng W, Chen S, Tu M, Wu M, et al. Analysis of the Risk Factors for Mortality in Adult COVID-19 Patients in Wuhan: A Multicenter Study. *Front Med* [Internet]. 2020 [cited 2020 Nov 16];7. Available from: <https://www.frontiersin.org/articles/10.3389/fmed.2020.00545/full>
33. Albitar O, Ballouze R, Ooi JP, Sheikh Ghadzi SM. Risk factors for mortality among COVID-19 patients. *Diabetes Res Clin Pract*. 2020 Aug 1;166:108293.
34. Holman N, Knighton P, Kar P, O'Keefe J, Curley M, Weaver A, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol*. 2020 Oct 1;8(10):823–33.
35. Zheng Yang, Qinming Hu, Fei Huang, Shouxin Xiong, Yi Sun. The prognostic value of the SOFA score in patients with COVID-19: a retrospective, observational study. 2020 Oct 26 [cited 2020 Nov 20]; Available from: <https://europepmc.org/article/ppr/ppr231113>

PLACID TRIAL Mortality Assessment

Table 1: Distribution of demographic variables and clinical parameters in enrolled patients and comparison of survivors and non-survivors in the cohort

Variables		Overall (n=451) N (%)	Survivor (n=386) N (%)	Non survivor (n=65) N (%)	P value
Age (Mean ± SD)		51±12.4	50±12.4	56±11.3	<0.001*
Age	≤ 40	104 (23.1)	97 (25.1)	7 (10.8)	0.004
	41-59	225 (49.9)	194 (50.3)	31 (47.7)	
	≥60	122 (27.1)	95 (24.6)	27 (41.5)	
Gender: Male		346 (76.7)	294(76.2)	52 (80.0)	0.499
Blood group	A	104(23.1)	91(23.6)	13(20.0)	0.518
	B	164(36.4)	140(36.3)	24(36.9)	
	AB	25(5.5)	19(4.9)	6(9.2)	
	O	158(35.0)	136(35.2)	22(33.8)	
History of smoking		37(8.2)	32(8.3)	5(7.7)	0.866
Comorbidities and Chronic illness					
Diabetes		196 (43.5)	164 (42.5)	32 (49.2)	0.310
Hypertension		169 (37.5)	139 (36.0)	30 (46.2)	0.118
Chronic obstructive pulmonary disease		15 (3.3)	10 (2.6)	5 (7.7)	0.050
Obesity ≥ 30		31 (6.9)	25 (6.5)	6 (9.2)	0.426
Chronic kidney disease		17 (3.8)	11 (2.8)	6 (9.2)	0.024
Coronary artery disease		31 (6.9)	23 (6.0)	8 (12.3)	0.106
Cerebrovascular disease		4 (0.9)	3 (0.8)	1 (1.5)	0.465
Symptoms at admission					
Shortness of breath		413 (91.6)	351 (90.9)	62 (95.4)	0.232
Fever		158 (35.0)	128 (33.2)	30 (46.2)	0.042
Cough		309 (68.5)	259 (67.1)	50 (76.9)	0.115
Fatigue		354 (78.7)	301 (78.2)	53 (81.5)	0.541
Severity of illness score					
SOFA score at admission*		2.40 ± 1.06	2.30±0.93	3.05±1.49	<0.001
Treatment					
Vasopressor		18 (4.0)	1 (0.3)	17 (26.6)	<0.001
Non-Invasive Ventilation (NIV)		446 (98.9)	383 (99.2)	63 (96.9)	0.101
Invasive ventilation		38 (8.4)	4 (1.04)	34 (52.31)	<0.001
Interval between symptoms onset to admission ‡		4 (3, 7)	4 (3, 7)	4 (3, 6)	0.996
Duration of respiratory support days ‡		6 (4, 10)	6 (4, 9.5)	6 (3, 10)	0.689
Duration of invasive ventilation days ‡		1 (1,3)	12 (2, 14)	1 (1, 3)	0.020
Duration of hospital stay days ‡		14 (10, 18)	14 (11, 19)	8 (5, 14)	<0.001

‡ Median (IQR) days in days – Mann Whitney U test was used

*Mean ± SD – Independent t test was used

PLACID TRIAL Mortality Assessment

Table 2: Univariate Fine and Gray model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Univariate Analysis					
		Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00		
	41-59	2.04	0.90 - 4.66	0.089	0.80	0.64 - 1.01	0.057
	≥60	3.51	1.53 - 8.07	0.003	0.56	0.42 - 0.73	<0.001
Gender	Male	1.19	0.64 - 2.19	0.582	0.87	0.70 - 1.09	0.228
Blood Group	O	1.00			1.00		
	A	0.94	0.48 - 1.87	0.866	0.89	0.69 - 1.15	0.389
	B	1.13	0.63 - 2.01	0.689	0.93	0.75 - 1.18	0.578
	AB	2.01	0.80 - 5.05	0.139	0.66	0.39 - 1.11	0.116
Comorbidities	No Comorbidities	1.00			1.00		
	1	1.62	0.82 - 3.21	0.166	0.79	0.63 - 0.99	0.044
	2 or More	2.25	1.18 - 4.29	0.014	0.70	0.55 - 0.88	0.003
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00		
	5-10	4.90	1.80 - 13.32	0.002	0.72	0.56 - 0.93	0.013
	>10	28.84	11.92 - 69.76	<0.001	0.17	0.12 - 0.26	<0.001
Platelet count † (* 10 ⁹ /L)	<100	6.88	3.61 - 13.13	<0.001	0.16	0.05 - 0.49	0.001
	≥ 100	1.00			1.00		
SOFA score †		1.63	1.54 - 1.74	<0.001	0.62	0.57 - 0.67	<0.001
D-dimer(mg/L) \$	<0.5	1.00			1.00		
	0.5 - 1.0	1.53	0.63 - 3.67	0.346	0.82	0.64 - 1.06	0.129
	>1.0	3.34	1.55 - 7.19	0.002	0.57	0.45 - 0.73	<0.001
Ferritin(ng/mL) \$	<500	1.00			1.00		
	≥500	4.11	2.28 - 7.41	<0.001	0.52	0.42 - 0.64	<0.001
CRP ^s (mg/L)		1.0003	0.999 - 1.001	0.080	0.99	0.99 - 1.00	0.360
LDH ^s (IU/L)	<450	1.00			1.00		
	≥ 450	4.88	2.72 - 8.75	<0.001	0.53	0.43 - 0.66	<0.001
PaO ₂ /FiO ₂ ‡	<100 (severe)	25.64	14.8 - 44.41	<0.001	6.5e-08	4.3e-08 - 9.9e-08	<0.001
	100-200(moderate)	5.97	3.05 - 11.69	<0.001	0.19	0.10 - 0.36	<0.001
	>200 (Mild)	1.00			1.00		
Interval from onset of symptoms to admission		1.05	0.96 - 1.14	0.334	0.97	0.94 - 1.00	0.058
Vasopressor support		11.36	7.79 - 16.56	<0.001	0.03	0.004 - 0.22	0.001
Invasive ventilation support		19.57	12.21 - 31.35	<0.001	0.01	0.002 - 0.09	<0.001

† Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: \$ inflammatory biomarkers values were measured at day 0, 3 and day 7.SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

PLACID TRIAL Mortality Assessment

478

479 Table 3: Multivariable Fine and Gray Model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Multivariable Analysis (Model A)						Multivariable Analysis (Model B)					
		Mortality			Discharged alive			Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00			1.00			1.00		
	41-59	1.23	0.60 - 2.49	0.572	1.03	0.81 - 1.32	0.825	1.55	0.65 - 3.71	0.325	0.95	0.74 - 1.21	0.671
	≥60	1.44	0.67 - 3.09	0.347	0.94	0.70 - 1.26	0.675	1.72	0.67 - 4.46	0.261	0.78	0.57 - 1.06	0.110
Comorbidities	No Comorbidities	1.00			1.00			1.00			1.00		
	1	1.20	0.69 - 2.10	0.515	0.91	0.72 - 1.14	0.407	1.31	0.59 - 2.91	0.509	0.87	0.67 - 1.11	0.254
	2 or More	1.76	1.02 - 3.03	0.041	0.89	0.69 - 1.14	0.329	2.68	1.26 - 5.70	0.011	0.69	0.52 - 0.93	0.014
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00								
	5-10	3.34	1.21 - 9.19	0.020	0.81	0.64 - 1.03	0.095						
	>10	9.97	3.65 - 27.13	<0.001	0.39	0.26 - 0.58	<0.001						
SOFA score †		1.22	1.11 - 1.35	<0.001	0.75	0.68 - 0.83	<0.001						
D-dimer(mg/L) §	<0.5							1.00					
	0.5 - 1.0							1.29	0.54 - 3.10	0.565	0.84	0.65 - 1.09	0.198
	>1.0							2.50	1.14 - 5.48	0.022	0.64	0.49 - 0.82	<0.001
Ferritin(ng/mL) §	<500							1.00			1.00		
	≥500							2.67	1.44 - 4.96	0.002	0.69	0.55 - 0.86	0.001
LDH§ (IU/L)	<450							1.00			1.00		
	≥ 450							2.96	1.60 - 5.45	0.001	0.68	0.55 - 0.85	0.001
PaO2/FiO2‡	<100 (severe)	3.47	1.64 - 7.37	0.001	3.1e-07	1.7e-07 - 5.7e-07	<0.001						
	100-200(moderate)	1.91	0.91 - 4.004	0.087	0.401	0.19 - 0.85	0.016						
	>200 (Mild)	1.00			1.00								

480

481 †Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: § Inflammatory biomarkers values were measured at day 0, 3 and day 7

482 (Model A) Multivariable Fine and Gray model for Age, comorbidities with Laboratory Parameters, (Model B) Multivariable Fine and Gray model for Age, comorbidities with
483 inflammatory biomarker values

484 SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

485

486

PLACID TRIAL Mortality Assessment

Figure 1:





Figure 1: Line graph for SOFA score showing difference between survivors and non survivors

Figure 1 showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

PLACID TRIAL Mortality Assessment

Supplementary

Flowchart for the study protocol



The Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID) Trial recruited 464 eligible patients for the study. The primary outcome at 28 days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomization and two patients did not receive the intervention after randomization as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients out of whom 386 survived and 65 died.

BMJ Open

Factors associated with mortality among moderate and severe COVID 19 patients in India – A secondary analysis of a Randomised Controlled Trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-050571.R2
Article Type:	Original research
Date Submitted by the Author:	31-Jul-2021
Complete List of Authors:	<p>Mammen, Joy; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Kumar, Snehil; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Thomas, Lovely; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Kumar, Gunjan ; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Zachariah, Anand; Christian Medical College and Hospital Vellore, Medicine</p> <p>Jeyaseelan, Lakshmanan; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Peter, John Victor; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Agarwal, Anup; ICMR, Clinical Trial and Health Systems Research Unit</p> <p>Mukherjee, Aparna; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Chatterjee, Pranab; ICMR, Translational Global Health Policy and Research Cell</p> <p>Bhatnagar, Tarun ; National Institute of Epidemiology, ICMR School of Public Health</p> <p>Rasalam, Jess; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Chacko, Binila; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Mani, Thenmozhi; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Joy, Melvin; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Rupali, Priscilla; Christian Medical College and Hospital Vellore, Infectious Diseases</p> <p>Murugesan, Malathi; Christian Medical College and Hospital Vellore, Hospital Infection Control Committee</p> <p>Daniel, Dolly; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Latha, B; Madras Medical College, Transfusion Medicine</p> <p>Bundas, Sunita; SMS Medical College and Hospital, Transfusion Medicine</p> <p>Kumar, Vivek; Sir HN Reliance Foundation Hospital and Research Centre, Critical Care</p> <p>Dosi, Ravi; Sri Aurobindo Institute of Medical Sciences, Respiratory Medicine</p>

	Khambholja, Janakkumar; Smt NHL Municipal Medical College, Internal Medicine de Souza, Rosemarie; BYL Nair Charitable Hospital, Department of Medicine Chander, B; Gandhi Medical College and Hospital, Internal Medicine Bahadur, Shalini; Government Institute of Medical Sciences, Pathology Dube, Simmi; Gandhi Medical College Bhopal, Internal Medicine Suri, Amit; Atal Bihari Vajpayee Institute of Medical Sciences and Ram Manohar Lohia Hospital, Pulmonary Medicine Jindal, Aikaj ; Satguru Pratap Singh Hospitals, Transfusion Medicine Shrivastav, Om ; Kasturba Hospital for Infectious Diseases, Infectious diseases Barge, Vijay; RSCM Government Medical College, Medicine Bajpayee, Archana; AIIMS Jodhpur, Transfusion Medicine Malhotra, Pankaj; Post Graduate Institute of Medical Education and Research, Department of Internal Medicine Singh, Neha; AIIMS Patna, Transfusion Medicine Tambe, Muralidhar; Department of Community Medicine, B J Govt. Medical College Sharma, Nimisha; ESIC Medical College and Hospital Faridabad, Transfusion Medicine Bhat, Shreepad; Smt Kashibai Navale Medical College and General Hospital, Internal Medicine Kaulgud, Ram; Karnataka Institute of Medical Sciences, Internal Medicine Gurtoo, Anil; Lady Hardinge Medical College, Internal Medicine Reddy, Himanshu; King George Medical College, Internal Medicine Upadhyay, Kamlesh; Byramjee Jeejabhoy Medical College, Internal medicine Jain, Ashish ; Mahatma Gandhi Medical College and Hospital, Respiratory Medicine Patel, Tinkal; Government Medical College, Internal Medicine Nagori, Irfan; GMERS Medical College Gotri Vadodara, Medicine Jha, Pramod ; Sumandeep Vidyapeeth University, Internal Medicine Babu, KV; Sri Venkateswara Institute of Medical Sciences, Transfusion Medicine Aparna, C; Kurnool Medical College, Pathology Panjwani, Sunil; Government Medical College, Internal Medicine Natarajan, M; Madurai Medical College, Internal Medicine Baldi, Milind; Mahatama Gandhi Memorial Medical College, Internal Medicine Khadke, Vrushali; Poona Hospital and Research Centre, Interventional Pulmonology Dua, Seema; Super Speciality Paediatric Hospital and Teaching Hospital, Transfusion Medicine Singh, Ravindraa; Aditya Birla Memorial Hospital, Transfusion Medicine Sharma, Ashish; R. D. Gardi Medical College, Medicine Sharma, Jayashree; Seth GS Medical College and KEM Hospital, Transfusion Medicine Gokhale, Yojana; Lokmanya Tilak Municipal Medical College and General Hospital, Internal Medicine Yadav, Pragya; National Institute of Virology, Maximum Containment Laboratory, ICMR Sapkal, Gajanan; National Institute of Virology, Diagnostic Virology Group, ICMR Kaushal, Himanshu; National Institute of Virology, Human Influenza Group, ICMR Kumar, Saravana; National Institute of Epidemiology, Epidemiology and Biostatistics Division
Primary Subject	Public health

Heading:	
Secondary Subject Heading:	Public health
Keywords:	COVID-19, Public health < INFECTIOUS DISEASES, Adult intensive & critical care < INTENSIVE & CRITICAL CARE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

PLACID TRIAL Mortality Assessment**Factors associated with mortality among moderate and severe COVID 19 patients in India – A secondary analysis of a Randomised Controlled Trial**

Joy J Mammen, Snehil Kumar, Lovely Thomas, Gunjan Kumar, Anand Zachariah, Lakshmanan Jeyaseelan, John Victor Peter, Anup Agarwal, Aparna Mukherjee, Pranab Chatterjee, Tarun Bhatnagar, Jess Elizabeth Rasalam, Binila Chacko, Thenmozhi Mani, Melvin Joy, Priscilla Rupali, Malathi Murugesan, Dolly Daniel, B Latha , Sunita Bundas, Vivek Kumar, Ravi Dosi, Janakkumar R Khambholja, Rosemarie de Souza, B Thrilok Chander, Shalini Bahadur, Simmi Dube, Amit Suri, Aikaj Jindal, Om Shrivastav, Vijay Barge, Archana Bajpayee, Pankaj Malhotra, Neha Singh, Muralidhar Tambe, Nimisha Sharma, Shreepad Bhat, Ram S Kaulgud, Anil Gurtoo, D Himanshu Reddy, Kamlesh Upadhyay, Ashish Jain, Tinkal C Patel, Irfan Nagori, Pramod R Jha, KV Sreedhar Babu, C Aparna, Sunil Jodharam Panjwani, M Natarajan, Milind Baldi, Vrushali Khirid Khadke, Seema Dua, Ravindraa Singh, Ashish Sharma, Jayashree Sharma, Yojana A Gokhale, Pragya D Yadav, Gajanan Sapkal, Himanshu Kaushal, V Saravana Kumar

Corresponding Author:

Joy J Mammen MD

Professor & Head

Address

Department of Transfusion Medicine and Immunohaematology

Christian Medical College Vellore

Tamil Nadu, India

Pin code 632004

E mail joymammen@cmcvellore.ac.in

Phone +91 416 228 2536

Fax +91 416 223 2035

PLACID TRIAL Mortality Assessment

32

Name	Designation and affiliation	Full Address
Joy J Mammen	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Snehil Kumar	Senior Resident, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lovely Thomas	Assistant Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Gunjan Kumar	Scientist C, Clinical Trial & Health Systems Research Unit, Indian Council of Medical Research (ICMR)	New Delhi, India.
Anand Zachariah	Professor, Department of Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lakshmanan Jeyaseelan	Professor, Clinical Epidemiology Unit, Christian Medical College	Vellore, Tamil Nadu, India.
John Victor Peter	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Anup Agarwal	Consultant, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Aparna Mukherjee	Scientist E, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Pranab Chatterjee	Scientist B, Translational Global Health Policy and Research Cell, ICMR	New Delhi, India.
Tarun Bhatnagar	Scientist E, ICMR School of Public Health, ICMR -National Institute of Epidemiology	Chennai, Tamil, India
Jess Elizabeth Rasalam	Fellow, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Binila Chacko	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Thenmozhi Mani	Senior Demonstrator, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Melvin Joy	Associate Research Officer, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Priscilla Rupali	Professor, Department of Infectious Diseases, Christian Medical College	Vellore, Tamil Nadu, India.
Malathi Murugesan	Assistant Professor, Hospital Infection Control Committee, Christian Medical College	Vellore, Tamil Nadu, India.
Dolly Daniel	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
B Latha	Professor, Department of Transfusion Medicine, Madras Medical College	Chennai, Tamil Nadu, India
Sunita Bundas	Professor, Department of Immunohaematology and Transfusion Medicine, SMS Medical College and Hospital	Jaipur, Rajasthan, India
Vivek Kumar	Professor, Department of Critical Care Medicine, Sir H N Reliance Foundation Hospital	Mumbai, Maharashtra, India
Ravi Dosi	Professor, Department of Respiratory Medicine, Sri Aurobindo Institute of Medical Sciences	Indore, Madhya Pradesh, India
Janakkumar R Khambholja	Professor, Department of Internal Medicine, Smt NHL Municipal Medical College	Ahmedabad, Gujarat, India
Rosemarie de Souza	Professor, Department of Internal Medicine, BYL Nair Charitable Hospital & TN Medical College	Mumbai, Maharashtra, India
B Thrilok Chander	Professor, Department of Internal Medicine, Gandhi Medical College, Secunderabad	Secunderabad, Telangana, India
Shalini Bahadur	Associate Professor, Department of Pathology, Government Institute of Medical Sciences	Greater Noida, Uttar Pradesh, India
Simmi Dube	Professor, Department of Internal Medicine, Gandhi Medical College	Bhopal, Madhya Pradesh, India
Amit Suri	Professor, Department of Pulmonary Medicine, Atal Bihari Vajpayee Institute of Medical Sciences and RML Hospital	New Delhi, India
Aikaj Jindal	Professor, Department of Transfusion Medicine, Satguru Partap Singh Hospitals	Ludhiana, Punjab, India
Om Shrivastav	Professor, Department of Infectious Diseases, Kasturba Hospital	Mumbai, Maharashtra, India
Vijay Barge	Professor, Department of Medicine, RSCM Government Medical College	Kolhapur, Maharashtra,
Archana Bajpayee	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Jodhpur, Rajasthan, India
Pankaj Malhotra	Professor, Department of Internal Medicine, Post Graduate Institute of Medical Education and Research	Chandigarh, India
Neha Singh	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Patna, Bihar, India
Muralidhar Tambe	Professor, Department of Community Medicine Byramjee Jeejeebhoy Medical College	Pune, Maharashtra, India

PLACID TRIAL Mortality Assessment

Nimisha Sharma	Professor, Department of Immunohematology and blood transfusion, ESIC Medical College and Hospital	Faridabad, Haryana, India
Shreepad Bhat	Professor, Department of Internal Medicine, Smt. Kashi Bai Navale Medical College	Pune, Maharashtra, India
Ram S Kaulgud	Professor, Department of Internal Medicine, Karnataka Institute of Medical Sciences	Hubali, Karnataka, India
Anil Gurtoo	Professor, Department of Internal Medicine, Lady Hardinge Medical College	New Delhi, India
D Himanshu Reddy	Professor, Department of Internal Medicine, King George's Medical University	Lucknow, Uttar Pradesh, India
Kamlesh Upadhyay	Professor, Department of Internal Medicine, Byramjee Jeejeebhoy Medical College	Ahmedabad, Gujarat, India
Ashish Jain	Professor, Department of Respiratory Medicine, Mahatma Gandhi University of Medical Sciences and Technology	Jaipur, Rajasthan, India
Tinkal C Patel	Professor Department of Internal Medicine, Government Medical College	Surat, Gujarat, India
Irfan Nagori	Professor, Department of Medicine, GMERS Medical College and Hospital	Vadodara, Gujarat, India
Pramod R Jha	Professor, Department of Internal Medicine, Sumandeep, Vidyapeeth and Dhiraj Hospital	Vadodara, Gujarat, India
KV Sreedhar Babu	Professor, Department of Transfusion Medicine, Sri Venkateswra Institute of Medical Sciences	Tirupati, Andhra Pradesh, India
C Aparna	Professor, Department of Pathology, Kurnool Medical College	Kurnool, Andhra Pradesh, India
Sunil Jodharam Panjwani	Professor, Department of Internal Medicine, Government Medical College	Bhavnagar, Gujarat, India
M Natarajan	Professor, Department of Internal Medicine, Madurai Medical College	Madurai, Tamil Nadu, India
Milind Baldi	Professor, Department of Respiratory Medicine, Mahatma Gandhi Memorial Medical College	Indore, Madhya Pradesh, India
Vrushali Khirid Khadke	Professor, Department of Interventional Pulmonology, Poona Hospital and Research Center	Pune, Maharashtra, India
Seema Dua	Professor, Department of Transfusion Medicine, Super Specialty Paediatric Hospital and Post Graduate Teaching Institute	Noida, Uttar Pradesh, India
Ravindraa Singh	Professor, Department of Transfusion Medicine, Aditya Birla Memorial Hospital	Pune, Maharashtra, India
Ashish Sharma	Professor, Department of Medicine, RD Gardi Medical College	Ujjain, Madhya Pradesh, India
Jayashree Sharma	Professor, Department Transfusion Medicine, Seth GS Medical College and KEM Hospital	Mumbai, Maharashtra, India
Yojana A Gokhale	Professor, Department of Internal Medicine, Lokmanya Tilak Municipal Medical College & General Hospital	Mumbai, Maharashtra, India
Pragya D Yadav	Scientist E, Maximum Containment Laboratory, ICMR – National Institute Virology	Pune, Maharashtra, India
Gajanan Sapkal	Scientist E, Diagnostic Virology Group, ICMR – National Institute Virology	Pune, Maharashtra, India
Himanshu Kaushal	Scientist B, Human Influenza group, ICMR – National Institute Virology	Pune, Maharashtra, India
V Saravana Kumar	Scientist B, Epidemiology and Biostatistics Division, ICMR – National Institute of Epidemiology	Chennai, Tamil Nadu, India

33

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

PLACID TRIAL Mortality Assessment

Abstract

Objective:

Large data on the clinical characteristics and outcome of COVID-19 in the Indian population is scarce. We analysed the factors associated with mortality in a cohort of moderate and severely ill COVID-19 patients enrolled in a randomised trial on convalescent plasma.

Design:

Secondary analysis of data from a Phase II, Open Label, Randomised Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID TRIAL).

Setting:

39 public and private hospitals across India, during the study period 22 April 2020 to 14 July 2020.

Participants:

Of the 464 patients recruited, two were lost to follow-up, nine withdrew consent and two patients did not receive the intervention after randomisation. The cohort of 451 participants with known outcome at 28-days was analysed.

Primary outcome measure:

Factors associated with all-cause mortality at 28-days post-enrolment.

Results:

The mean (SD) age was 51±12.4 years; 76.7% were males. Admission SOFA score was 2.4±1.1. Non-invasive ventilation, invasive ventilation and vasopressor therapy were required in 98.9%, 8.4% and 4.0% respectively. The 28-day mortality was 14.4%. Median time from symptom onset to hospital admission was similar in survivors (4 days; IQR 3-7) and non survivors (4 days; IQR 3-6). Patients with two or more co-morbidities had 2.25 (95%CI: 1.18-4.29, p=0.014) times risk of death. When compared with survivors, admission IL-6 levels were higher (p<0.001) in non-survivors and increased further on Day 3. On multivariable Fine and Gray model , severity of illness (SHR 1.22, 95%CI:1.11-1.35,p<0.001), PaO₂/FiO₂ ratio <100 (3.47, 1.64-7.37, p=0.001), Neutrophil Lymphocyte ratio (NLR) >10 (9.97, 3.65-27.13, p<0.001), D-dimer >1.0mg/L (2.50,1.14-5.48, p=0.022), ferritin ≥500ng/mL (2.67, 1.44-4.96, p=0.002) and LDH ≥450 IU/L (2.96, 1.60-5.45, p=0.001) were significantly associated with death.

Conclusion:

In this cohort of moderate and severely ill COVID-19 patients, severity of illness, underlying co-morbidities and higher levels of inflammatory markers were significantly associated with death.

PLACID TRIAL Mortality Assessment**Trial Registration:**

CTRI/2020/04/024775.

For peer review only

1		PLACID TRIAL Mortality Assessment
2		
3		
4	90	Article Summary
5		
6	91	Strengths and limitations of this study
7		
8		
9	92	Strengths
10		
11	93	There is no study from India, with representation from multiple states that has detailed the
12		
13	94	clinical profile and evaluated for factors associated with death. This study may help with strategic
14		
15	95	planning at a national level.
16		
17	96	The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28,
18		
19	97	did not differ across the trial arms, therefore the present analysis need not be adjusted for
20		
21	98	convalescent plasma intervention.
22		
23		
24	99	There may be variability of treatment provided in the multiple centres, however, care was taken
25		
26	100	that patients received best standard of care for COVID-19 dictated by the best available evidence at
27		
28	101	the time and guidelines for the management of COVID-19 issued by health authorities of the Indian
29	102	government.
30		
31		
32	103	Limitations
33		
34	104	The laboratory and biomarker assays for ferritin, lactate dehydrogenase, C reactive protein, and D-
35		
36	105	dimer were conducted using tests from different manufacturers.
37		
38		
39	106	Participants of this study may not comprise a true observational cohort as this was a post hoc
40		
41	107	analysis of a randomised control trial data. Our study did not analyse the effect of SARS-CoV-2
42		
43	108	variants causing a high mortality in younger population during the second wave of COVID-19
44	109	infection, and therefore extrapolation to the general population must be carefully qualified.
45		
46	110	
47		
48		
49	111	
50		
51		
52	112	
53		
54	113	
55		
56		
57		
58		
59		
60		

PLACID TRIAL Mortality Assessment**Introduction**

The first human case of Corona Virus Disease 19 (COVID-19) caused by the novel coronavirus (named Severe Acute Respiratory Syndrome Coronavirus 2, SARS-CoV-2) was reported in Wuhan City, China in December 2019. On 30 January 2020, the World Health Organisation (WHO) declared that the outbreak of COVID-19 constituted a Public Health Emergency of International Concern (1). Based on the high levels of global spread and the severity of COVID-19, on 11 March 2020, the Director-General of the WHO declared the COVID-19 outbreak a pandemic (2). The sudden outbreak followed by rapid spread in a globalised world, resulted in a major medical burden, besides affecting socio-economic well-being among all nations.

In India, the disease was first detected on 30 January 2020 in the state of Kerala, in a student who returned from Wuhan (3). After a brief, initial respite, the virus has spread at a rapid pace in India, resulting in more than 10 million confirmed cases as of December, 2020 with more than 145,000 deaths (4).

Patients diagnosed with COVID-19 have primarily respiratory symptoms. Most patients diagnosed with COVID-19 experience mild to moderate respiratory illness, fever, dry cough, fatigue and recover without requiring special treatment (5). Oxygen desaturation is the hallmark of progression. Patients with underlying medical problems like cardiovascular disease, diabetes, chronic respiratory disease, and cancer are more likely to develop serious illness. These patients may develop viral pneumonia, with resultant dyspnoea and hypoxaemia which may progress to respiratory or multi- system failure and even death (6). There is paucity of large-scale data on the clinical characteristics, outcomes of COVID-19 in the Indian population and evaluation of risk factors with an unfavourable outcome at a national level. Identification of such potential risk factors is important to anticipate medical treatment and to reduce the mortality burden for severe COVID-19 illness by proactive interventions.

The Indian Council of Medical Research (ICMR) conducted a randomised trial (A Phase II, Open Label, Randomised Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease, PLACID TRIAL) to determine the effectiveness and safety of convalescent plasma in patients with moderate and severely ill COVID-19 to limit progression of disease (7). Patients received standard of care for COVID-19 in keeping with the institutional protocols, based on the best available evidence at the time and guidelines for the management of COVID-19 issued by national health authorities. Participants in the intervention

PLACID TRIAL Mortality Assessment

arm received two doses of 200 mL of convalescent plasma, transfused 24 hours apart, in addition to standard of care. The control arm did not receive any additional therapy. The study concluded that the use of convalescent plasma was not associated with a reduction in 28-day mortality (7). The aim of this analysis was to identify risk factors associated with mortality by mining the data collected from the cohort enrolled in the PLACID TRIAL (7).

Methods

Participants

The study enrolled patients from 39 different hospitals, of which, 29 were teaching public hospitals and 10 were private facilities spread across 14 states and union territories. Patients over the age of 18 years who were confirmed to have COVID-19 based on a positive SARS-CoV-2 RT-PCR test and moderate and severely ill with either a partial pressure of oxygen in arterial blood/fraction of inspired oxygen (PaO₂/FiO₂) ratio between 200-300 or respiratory rate >24/min and decreased oxygen saturation on room air (SpO₂ < 93%) were included during the study period from 22 April to 14 July 2020. As per the guidelines issued by the Ministry of Health, Government of India at the time of conduct of the study, the subset of patients with the above criteria but with a respiratory rate between 24 and 30/min were classified as moderate disease. Those with respiratory rate >30 breaths/min were classified as severe disease(8). Patients were followed up for 28-days and assessed for their health status and all-cause mortality. Ethics approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals. Written consent was obtained from patients or their families before enrolling in the study.

Data

Data was obtained from the ICMR PLACID TRIAL database collected in structured paper case record forms and entered in Research Electronic Data Capture system (REDCap, version 8.5 Vanderbilt University, TN). The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775). After trial completion, based on co-operative agreement between the centres, and IRB permission, the data was shared and analysed further, to explore for other meaningful results. No separate ethical clearance was taken for this study.

PLACID TRIAL Mortality Assessment

Demographic, clinical, laboratory tests and outcome data were collected prospectively. Clinical symptoms need for organ support (respiratory, haemodynamic), laboratory tests (complete blood count, coagulation profile, serum biochemical profile, renal and liver function tests) were monitored serially on day of enrolment (day 0) and on day 1, 3, 5, 7, 14 and 28. Inflammatory biomarkers [lactate dehydrogenase (LDH), serum ferritin, and C-reactive protein (CRP)] were tested at admission and on day 3 and 7 whereas; Interleukin 6 (IL-6) was done at admission and on day 3.

The outcome of interest was all-cause day 28 mortality. We evaluated for association between laboratory parameters and mortality.

Statistical Methods:

Mean and standard deviation (SD) or median and inter-quartile range (IQR) were used for continuous variables as appropriate, and categorical variables number and proportions were used. To find the association between mortality and study variables, Chi-square test/Fisher's exact test were used. To find the mean difference across the groups, independent t test was used. Similarly, Mann Whitney U test was used to compare median difference. The end point of interest was all-cause mortality (event of interest) at day 28 from the time of enrolment, discharged alive (competing event) and hospital admission after day 28 (censored), whichever is earlier. Discharged alive was treated as a competing event because the event of discharged alive precludes the event of all-cause mortality. The variables that are statistically significant or clinically important are considered in the multivariable Fine and Gray regression model. However, if a variable is expected to have collinear concern or had sparse data that was not included in the analysis. Two multivariable models were developed. The first model included clinical and laboratory parameters tested on day 0, 1, 3, 5, 7, 14 and 28 while the second included inflammatory biomarkers tested on day 0, 3 and 7 after adjusting for age and comorbidities. For certain laboratory markers such as D-dimer, ferritin and LDH, clinically relevant thresholds were used for the analysis rather than using these data as continuous variables. The clinically relevant thresholds for these variables were set as >1.0 mg/L for D-dimer, ≥ 500 mg/mL for Ferritin and ≥ 450 IU/L for LDH. The threshold for Ferritin of 500 μ g/L was based on the cut-off value for the diagnosis of Hemophagocytic lymphohistiocytosis (HLH) as well as some preliminary evidence in COVID that a threshold of >500 μ g/L was associated with invasive ventilator dependence(9). Similarly, traditionally a threshold of <0.5 mg/L is used to exclude pulmonary thromboembolism; in this context two thresholds were used, 0.5 to 1.0 mg/L

PLACID TRIAL Mortality Assessment

and >1.0 mg/L(10). Variables included parameters that were strongly associated with mortality at univariate analysis or known from previous literature to be strongly associated with outcome. The model assumption was verified using log-log S (t) plots and Global test. A p-value of less than 0.05 levels was considered as statistically significant. All statistical analysis were performed using STATA version 16.0 (StataCorp. 2019. College Station, TX).

Patient and public involvement

Patients and the public were not involved in the design, conduct, reporting, or dissemination plans of our research. The study results will be disseminated to the study participants via their treating doctors.

Results

The PLACID Trial recruited 464 eligible patients for the study. The primary outcome at 28-days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomisation and two patients did not receive the intervention after randomisation as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients (*supplementary*).

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the analysis did not adjust for convalescent plasma intervention. The distribution of patients in intervention and control arms were 50.3% (n=227) and 49.7% (n=224), respectively. The mean (SD) age of the cohort was 51 ± 12.4 years; 76.7% were males. Table 1 shows distribution of demographic variables and clinical parameters in the study population.

The most common presenting symptoms were shortness of breath (91.6%), fatigue (78.7%), cough (68.5%) and fever (35%). Comorbidities were present in 59.9% of patients; 31.7% had any one comorbidities and 28.2% had two or more comorbidities. The most frequent comorbidities were diabetes (43.5%), hypertension (37.5%), obesity (6.9%) and Chronic Obstructive Pulmonary Disease (COPD) (3.3%). There was history of smoking in 8.2%. The time from onset of symptoms to admission was four days (IQR 3-7 days). Majority of the patients required non-invasive (98.9%) ventilatory support. The median duration of respiratory support was six days (IQR 4-10 days). In this cohort, 4% patients required vasopressor support. None of the patients required Extra Corporeal Membrane Oxygenation (ECMO) or dialysis support.

PLACID TRIAL Mortality Assessment

The all-cause mortality at 28-days was 14.4% (95%CI: 11.5-17.9, n=65). Median time from symptom onset to hospital admission was four days in survivors (IQR 3-7 days) and non survivors (IQR 3-6 days). The frequency of shortness of breath, cough and fatigue were similar in survivors and non survivors; however, the presence of fever at admission was significantly ($p=0.042$) associated with death (table 1). Other than COPD and CKD (chronic kidney disease), other comorbidities were not significantly associated with death (table 1). Admission Sequential Organ Failure Assessment (SOFA) score was higher in non survivors. The need for invasive mechanical ventilation, duration of invasive mechanical ventilation and vasopressor therapy were associated with death (table 1).

On univariate analysis (table 2), there was an association between increasing age and mortality. Patients with two or more comorbidities had a 2.25 (95%CI: 1.18-4.29, $p=0.014$) times increased chance of mortality. There was a strong mortality association for platelet count $<100 \times 10^9/L$ (SHR 6.88, 95%CI: 3.61-13.13, $p<0.001$), neutrophil lymphocyte ratio (NLR) >10 (28.84, 11.92-69.76, $p<0.001$), LDH ≥ 450 IU/L (4.88, 2.72-8.75, $p<0.001$), D-dimer >1 mg/L (3.34, 1.55-7.19, $p=0.002$) and ferritin ≥ 500 ng/mL (4.11, 2.28-7.41, $p<0.001$). Admission IL-6 levels were significantly ($p<0.001$) higher (76.00, 18.27-171.77) in non survivors than in survivors (18.51, 4.26-56.86). By day 3, IL-6 levels dropped to 11.6 (2.64-45.84) in survivors while it nearly doubled in non-survivors (140.35, 21.56-427.36). CRP did not show any statistical significance (1.0003, 0.999-1.001, $p=0.080$).

The need for invasive ventilation and vasopressors were associated with death (table 2). Increasing SOFA score was associated with mortality (1.63, 1.54-1.74, $p<0.001$). The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score progressively increased between the two groups over time (figure 1). Mortality proportionately also increased with lower PaO_2/FiO_2 values with sub-distribution hazard ratio of 25.64 (14.8-44.41, $p<0.001$) in the severe group as compared to the mild group.

Two models were run for multivariable Fine and Gray regression model over a period of time (table 3). Model A included age, comorbidities, PaO_2/FiO_2 , NLR and SOFA score. Model A revealed significant sub-distribution hazard ratios for PaO_2/FiO_2 ratio <100 (3.47, 1.64-7.37, $p=0.001$), NLR >10 (9.97, 3.65-27.13, $p<0.001$), SOFA score (1.22, 1.11- 1.35, $p<0.001$) after adjusting for age and comorbidities. Model B included age, comorbidities, D-dimer, ferritin and LDH. D-dimer >1 mg/L (2.50, 1.14-5.48, $p=0.022$), ferritin ≥ 500 ng/mL (2.67, 1.44-4.96, $p=0.002$) and LDH ≥ 450 IU/L

PLACID TRIAL Mortality Assessment

(2.96, 1.60-5.45, p=0.001), were associated with mortality after adjusting for age and comorbidities (table 3). IL-6 was omitted from the model as it was not measured on Day 7.

Discussion

In this study that enrolled patients from across India, we were able to identify clinical, biomarkers (D-dimer, LDH, ferritin) and SOFA score as factors that could indicate increased risk of death in moderate and severely ill COVID-19 patients from PLACID trial cohort. The definition of clinical grading of severity is different in India as compared to other countries (11–15). Mortality of critically ill COVID-19 patients varies significantly among the published case series, ranging from 16% to 78% (16–22). Similarly two studies from Wuhan, which included moderately as well as critically ill patients, have shown mortality rates of 3.77% and 14.14% (23,24). This wide variability can be explained by differences in the age of the population, distribution of risk factors, health system response across different countries, varied treatment protocols and follow-up. In a series of critically ill patients in China, 28-day ICU mortality was 61.5% (25). In a multicentric study from Italy, the mortality risk for patients without respiratory failure at admission was of 1% after 15 days while survival in patients with moderate-to-severe respiratory failure ($PaO_2/FiO_2 \leq 200$ mm Hg) at admission was only 56% at 15 days (26). The fatality rate reported in Europe and the United States of America is significantly higher than in China (27). Therefore, findings obtained in a specific country might not be automatically extrapolated and national cohorts must be studied.

In our study population, mortality increased with age, which is similar to the pattern observed in other countries affected by COVID-19. Age seems to affect the time from hospitalisation to death. Age-specific death rates was quite similar in studies from Asia, Europe and North America (28). South Korea, Italy, France, Germany, England and Wales, and Spain COVID-19 attributed mortality rates rise by about 12%/year while the United States and Wuhan, China showed a slower rate of increase about 9.5%/year of age (29). In a meta-analysis of 61, 11,583 subjects, 23.2% of patients were aged ≥ 80 years and showed an average mortality rate of 12.10%, the lowest being in China (3.1%) and the highest in the United Kingdom (20.8%) and New York State (20.99%). In the same study, highest mortality rate was observed in patients aged ≥ 80 years. The largest increase in mortality risk was observed in patients aged 60 to 69 years compared with those aged 50 to 59 years (odds ratio 3.13, 95% CI: 2.61-3.76) (30).

Presence of comorbidities significantly increases the death risk of COVID-19. A higher risk of mortality was seen in our patients who had CKD and COPD. Meta-analysis, including 1389 COVID-

PLACID TRIAL Mortality Assessment

19 patients with 19·7% having severe disease showed a significant association of CKD with severe COVID-19 with pooled odds ratio as 3·03 (31). Similarly, the estimated mortality risk in patients with COPD was three times of those without ($p<0\cdot05$) (32). We found that 43·5% of our patients had diabetes which is markedly higher as compared to patients from Korea which showed that 16·97% had diabetes mellitus (33). Our analysis showed that the presence of diabetes was not significantly different between survivors and non survivors (42·5% vs. 49·2%, $p=0\cdot310$), in contrast to the study from South Korea (33) which showed a much higher mortality among diabetic patients than in those without (20·0% vs. 4·8%). Hypertension and obesity were not significantly different among survivors and non-survivors in our study. However, the presence of two or more comorbidities was associated with mortality in our study.

Fine-Gray model identified prognostic markers for mortality, most notably age ≥ 60 years, $\text{PaO}_2/\text{FiO}_2$ ratio <100 , $\text{NLR} >10$, platelet count $<100 \times 10^9/\text{L}$, ferritin $>500\text{ng/mL}$, $\text{LDH} >450 \text{ IU/L}$ and D-dimer $>1\text{mg/L}$. Our study showed similar findings when compared with studies from Wuhan (34). Older age, leukocytosis, and high LDH level have been reported to be risk factors associated with in-hospital death in other studies also (35–37). IL-6 levels were significantly different in survivors and non-survivors at admission. By day 3 survivors had reducing IL-6 while it nearly doubled in non survivors.

Mortality was higher amongst patients requiring invasive mechanical ventilation (SHR 19·57, 12·21–31·35, $p<0\cdot001$) and those requiring vasopressors (SHR 11·36, 7·79–16·56, $p<0\cdot001$). However, the median duration of invasive ventilation for survivors was 12 days (2, 14) and that for non-survivors was one day (1, 3). These results suggest that patients with acute respiratory failure from COVID-19 may recover, even with severe disease requiring longer ventilation, and that probably the sickest patients die early reflecting lower duration of invasive ventilation in non survivors. Therefore, invasive ventilation should be timely and effectively provided.

In our study, the SOFA score was recognised as a valuable tool that could be used to prognosticate outcome of patients with COVID-19. Competing risk regression models showed that the increase in SOFA score was related to mortality, with a clearly divergent pattern between the two groups. Thus, an increasing SOFA score over time may be a factor that can be used to identify a subset of patients who may have an unfavourable outcome. Studies have shown that the SOFA

PLACID TRIAL Mortality Assessment

score could be used to evaluate severity and 60-day mortality of COVID-19 with the optimal cut-off score of 5 (38).

Limitations of the study include the variability of treatment provided in the multiple centres. The participants of this study may not comprise a true observational cohort as this was a post hoc analysis of a randomised control trial data and extrapolation to the general population must be carefully qualified. Our study did not analyse the effect of SARS-CoV-2 variants causing a high mortality in younger population during the second wave of COVID-19 infection and this may limit generalisability of the data to the second wave. Despite these limitations, this study provides a comprehensive overview of prognostic factors in moderate and severely ill COVID-19 patients that included patients from across the country.

Conclusion

Older age, multiple comorbidities, low PaO2/FiO2 ratio and deranged inflammatory markers are associated with worse prognosis. Serial SOFA score can be used for prognostication. Understanding symptoms, burden of comorbidities and systematic monitoring of key laboratory parameters offer opportunities for targeted intervention in COVID-19 with the use of anti-inflammatory or immunomodulatory agents.

Figure legend

Figure 1 showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

FOOTNOTES:

Authors Contributions:

Study design: JJM, LJ, JVP

Clinical Management: SK, LT, AZ, JER, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG.

PLACID TRIAL Mortality Assessment

Data collection: JJM, SK, LT, AZ, AA, AM, GK, PC, TB, JER, PR, MM, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG, PDY, GS, HK, VSK.

Data Analysis: JJM, SK, LJ, TM, MJ, PR, MM, DD, JVP.

Data Interpretation: JJM, SK, AZ, LJ, JER, TM, MJ, DD, JVP

Manuscript writing: JJM, SK, LT, AZ, LJ, JER, BC, TM, MJ, PR, MM, DD, JVP, AA, AM, GK, HK, PC, TB

Study Administration: JJM, BC, JVP, AA, AM, GK, HK, PC, TB

Funding support: The funding source for the primary study was ICMR (RFC No. ECD/NTF/20/2020-21/Covid Dated 23.07.2020). However, they had no financial interest in the investigational product. No other author has any competing financial or non-financial interest.

Competing interests: 'None declared'. All authors have completed the ICMJE uniform disclosure form at www.icmje.org.

Ethical Approval: Ethical approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) based in the National Center for Disease Informatics and Research, Indian Council of Medical Research, Bengaluru, Karnataka, as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals.

Data availability statement: Data will be made available, upon request, and must be accompanied by a brief proposal outlining the analysis plan. A signed data access agreement might be needed to ensure data safety and compliance with national rules about data sharing.

Acknowledgement: We acknowledge ICMR for providing us the data collected during PLACID trial which was analysed in the present study.

References:

1. Mullen L, Potter C, Gostin LO, Cicero A, Nuzzo JB. An analysis of International Health Regulations Emergency Committees and Public Health Emergency of International Concern Designations. BMJ Glob Health. 2020 Jun 1;5(6):e002502.
2. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. Acta Bio-Medica Atenei Parm. 2020 19;91(1):157–60.

PLACID TRIAL Mortality Assessment

- 386 3. Perappadan BS. India's first coronavirus infection confirmed in Kerala. The Hindu [Internet]. 2020
387 Jan 30 [cited 2020 Nov 16]; Available from: [https://www.thehindu.com/news/national/indias-](https://www.thehindu.com/news/national/indias-first-coronavirus-infection-confirmed-in-kerala/article30691004.ece)
388 [first-coronavirus-infection-confirmed-in-kerala/article30691004.ece](https://www.thehindu.com/news/national/indias-first-coronavirus-infection-confirmed-in-kerala/article30691004.ece)
- 389 4. MoHFW | Home [Internet]. [cited 2021 Jan 3]. Available from: <https://www.mohfw.gov.in/>
- 390 5. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. Clin Immunol Orlando
391 Fla. 2020 Jun;215:108427.
- 392 6. Wu C, Chen X, Cai Y, Xia J 'an, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory
393 Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan,
394 China. JAMA Intern Med. 2020 01;180(7):934–43.
- 395 7. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P. Convalescent plasma
396 in the management of moderate covid-19 in adults in India: open label phase II multicentre
397 randomised controlled trial (PLACID Trial). BMJ [Internet]. 2020 Oct 22 [cited 2020 Nov 16];371.
398 Available from: <https://www.bmj.com/content/371/bmj.m3939>
- 399 8. Government of India, Ministry of Health and Family Welfare, Directorate General of Health
400 Services. CLINICAL MANAGEMENT PROTOCOL FOR COVID-19-Version 3. Government of India;
401 2020.
- 402 9. Qeadan F, Tingey B, Gu LY, Packard AH, Erdei E, Saeed AI. Prognostic Values of Serum Ferritin
403 and D-Dimer Trajectory in Patients with COVID-19. Viruses. 2021 05;13(3).
- 404 10. Kearon C, de Wit K, Parpia S, Schulman S, Afilalo M, Hirsch A, et al. Diagnosis of Pulmonary
405 Embolism with d-Dimer Adjusted to Clinical Probability. N Engl J Med. 2019 Nov
406 28;381(22):2125–34.
- 407 11. Varghese GM, John R, Manesh A, Karthik R, Abraham OC. Clinical management of COVID-19.
408 Indian J Med Res. 2020 May 1;151(5):401.
- 409 12. Clinical management of COVID-19 [Internet]. [cited 2021 Jan 3]. Available from:
410 <https://www.who.int/publications/i/item/clinical-management-of-covid-19>
- 411 13. Zhang S-Y, Lian J-S, Hu J-H, Zhang X-L, Lu Y-F, Cai H, et al. Clinical characteristics of different
412 subtypes and risk factors for the severity of illness in patients with COVID-19 in Zhejiang, China.
413 Infect Dis Poverty. 2020 Jul 8;9(1):85.
- 414 14. Xia L, Chen J, Friedemann T, Yang Z, Ling Y, Liu X, et al. The Course of Mild and Moderate COVID-
415 19 Infections—The Unexpected Long-Lasting Challenge. Open Forum Infect Dis [Internet]. 2020
416 Sep 1;7(ofaa286). Available from: <https://doi.org/10.1093/ofid/ofaa286>
- 417 15. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of
418 coronavirus disease 2019: a model-based analysis. Lancet Infect Dis. 2020 Jun 1;20(6):669–77.
- 419 16. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al. Risk Factors Associated
420 With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy. JAMA
421 Intern Med. 2020 01;180(10):1345–55.
- 422 17. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019
423 novel coronavirus in Wuhan, China. Lancet Lond Engl. 2020 15;395(10223):497–506.

PLACID TRIAL Mortality Assessment

- 424 18. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, et al. Clinical Course and Outcomes of 344 Intensive
425 Care Patients with COVID-19. *Am J Respir Crit Care Med*. 2020 Apr 8;201(11):1430–4.
- 426 19. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized
427 Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA*. 2020 Mar
428 17;323(11):1061–9.
- 429 20. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, et al. Covid-19 in Critically Ill
430 Patients in the Seattle Region - Case Series. *N Engl J Med*. 2020 21;382(21):2012–22.
- 431 21. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and Outcomes
432 of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA*. 2020 28;323(16):1612–4.
- 433 22. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult
434 inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet Lond Engl*. 2020
435 28;395(10229):1054–62.
- 436 23. Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement,
437 and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect*. 2020 Jun 1;26(6):767–
438 72.
- 439 24. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased
440 patients with coronavirus disease 2019: retrospective study. *BMJ*. 2020 Mar 26;368:m1091.
- 441 25. Yang X, Yu Y, Xu J, Shu H, Xia J 'an, Liu H, et al. Clinical course and outcomes of critically ill
442 patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective,
443 observational study. *Lancet Respir Med*. 2020 May 1;8(5):475–81.
- 444 26. Santus P, Radovanovic D, Saderi L, Marino P, Cogliati C, De Filippis G, et al. Severity of respiratory
445 failure at admission and in-hospital mortality in patients with COVID-19: a prospective
446 observational multicentre study. *BMJ Open [Internet]*. 2020 Oct 10;10(10). Available from:
447 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7549463/>
- 448 27. Yamamoto N, Bauer G. Apparent difference in fatalities between Central Europe and East Asia
449 due to SARS-COV-2 and COVID-19: Four hypotheses for possible explanation. *Med Hypotheses*.
450 2020 Nov;144:110160.
- 451 28. Demographics of COVID-19 Deaths [Internet]. Ined - Institut national d'études démographiques.
452 [cited 2020 Nov 16]. Available from: <https://dc-covid.site.ined.fr/en/>
- 453 29. Goldstein JR, Lee RD. Demographic perspectives on the mortality of COVID-19 and other
454 epidemics. *Proc Natl Acad Sci U S A*. 2020 Sep 8;117(36):22035–41.
- 455 30. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et
456 al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583
457 Subjects. *J Am Med Dir Assoc*. 2020 Jul 1;21(7):915–8.
- 458 31. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019
459 (COVID-19) infection. *Int Urol Nephrol*. 2020 Mar 28;1–2.
- 460 32. Venkata VS, Kiernan G. COVID-19 AND COPD: POOLED ANALYSIS OF OBSERVATIONAL STUDIES.
461 *CHEST*. 2020 Oct 1;158(4):A2469.

PLACID TRIAL Mortality Assessment

33. D A, K L, Ds L, Ys L, Ss M. Mortality Rate and Predictors of Mortality in Hospitalized COVID-19 Patients with Diabetes. Healthc Basel Switz [Internet]. 2020 Sep 13 [cited 2020 Nov 16];8(3). Available from: <https://europepmc.org/article/med/32933191>

34. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. Eur Respir J [Internet]. 2020 May 7;55(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144257/>

35. Li M, Cheng B, Zeng W, Chen S, Tu M, Wu M, et al. Analysis of the Risk Factors for Mortality in Adult COVID-19 Patients in Wuhan: A Multicenter Study. Front Med [Internet]. 2020 [cited 2020 Nov 16];7. Available from: <https://www.frontiersin.org/articles/10.3389/fmed.2020.00545/full>

36. Albitar O, Ballouze R, Ooi JP, Sheikh Ghadzi SM. Risk factors for mortality among COVID-19 patients. Diabetes Res Clin Pract. 2020 Aug 1;166:108293.

37. Holman N, Knighton P, Kar P, O’Keefe J, Curley M, Weaver A, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. Lancet Diabetes Endocrinol. 2020 Oct 1;8(10):823–33.

38. Zheng Yang, Qinming Hu, Fei Huang, Shouxin Xiong, Yi Sun. The prognostic value of the SOFA score in patients with COVID-19: a retrospective, observational study. 2020 Oct 26 [cited 2020 Nov 20]; Available from: <https://europepmc.org/article/ppr/ppr231113>

PLACID TRIAL Mortality Assessment

Table 1: Distribution of demographic variables and clinical parameters in enrolled patients and comparison of survivors and non-survivors in the cohort

Variables	Overall (n=451) N (%)	Survivor (n=386) N (%)	Non survivor (n=65) N (%)	P value
Age (Mean ± SD)	51±12.4	50±12.4	56±11.3	<0.001*
Age				
≤ 40	104 (23.1)	97 (25.1)	7 (10.8)	
41-59	225 (49.9)	194 (50.3)	31 (47.7)	
≥60	122 (27.1)	95 (24.6)	27 (41.5)	0.004
Gender: Male	346 (76.7)	294(76.2)	52 (80.0)	0.499
Blood group				
A	104(23.1)	91(23.6)	13(20.0)	
B	164(36.4)	140(36.3)	24(36.9)	
AB	25(5.5)	19(4.9)	6(9.2)	
O	158(35.0)	136(35.2)	22(33.8)	0.518
History of smoking	37(8.2)	32(8.3)	5(7.7)	0.866
Comorbidities and Chronic illness				
Diabetes	196 (43.5)	164 (42.5)	32 (49.2)	0.310
Hypertension	169 (37.5)	139 (36.0)	30 (46.2)	0.118
Chronic obstructive pulmonary disease	15 (3.3)	10 (2.6)	5 (7.7)	0.050
Obesity ≥ 30	31 (6.9)	25 (6.5)	6 (9.2)	0.426
Chronic kidney disease	17 (3.8)	11 (2.8)	6 (9.2)	0.024
Coronary artery disease	31 (6.9)	23 (6.0)	8 (12.3)	0.106
Cerebrovascular disease	4 (0.9)	3 (0.8)	1 (1.5)	0.465
Symptoms at admission				
Shortness of breath	413 (91.6)	351 (90.9)	62 (95.4)	0.232
Fever	158 (35.0)	128 (33.2)	30 (46.2)	0.042
Cough	309 (68.5)	259 (67.1)	50 (76.9)	0.115
Fatigue	354 (78.7)	301 (78.2)	53 (81.5)	0.541
Severity of illness score				
SOFA score at admission*	2.40 ± 1.06	2.30±0.93	3.05±1.49	<0.001
Treatment				
Vasopressor	18 (4.0)	1 (0.3)	17 (26.6)	<0.001
Non-Invasive Ventilation (NIV)	446 (98.9)	383 (99.2)	63 (96.9)	0.101
Invasive ventilation	38 (8.4)	4 (1.04)	34 (52.31)	<0.001
Interval between symptoms onset to admission ‡	4 (3, 7)	4 (3, 7)	4 (3, 6)	0.996
Duration of respiratory support days ‡	6 (4, 10)	6 (4, 9.5)	6 (3, 10)	0.689
Duration of invasive ventilation days ‡	1 (1,3)	12 (2, 14)	1 (1, 3)	0.020
Duration of hospital stay days ‡	14 (10, 18)	14 (11, 19)	8 (5, 14)	<0.001

‡ Median (IQR) days in days – Mann Whitney U test was used

*Mean ± SD – Independent t test was used

PLACID TRIAL Mortality Assessment

Table 2: Univariate Fine and Gray model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Univariate Analysis					
		Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00		
	41-59	2.04	0.90 - 4.66	0.089	0.80	0.64 - 1.01	0.057
	≥60	3.51	1.53 - 8.07	0.003	0.56	0.42 - 0.73	<0.001
Gender	Male	1.19	0.64 - 2.19	0.582	0.87	0.70 - 1.09	0.228
Blood Group	O	1.00			1.00		
	A	0.94	0.48 - 1.87	0.866	0.89	0.69 - 1.15	0.389
	B	1.13	0.63 - 2.01	0.689	0.93	0.75 - 1.18	0.578
	AB	2.01	0.80 - 5.05	0.139	0.66	0.39 - 1.11	0.116
Comorbidities	No Comorbidities	1.00			1.00		
	1	1.62	0.82 - 3.21	0.166	0.79	0.63 - 0.99	0.044
	2 or More	2.25	1.18 - 4.29	0.014	0.70	0.55 - 0.88	0.003
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00		
	5-10	4.90	1.80 - 13.32	0.002	0.72	0.56 - 0.93	0.013
	>10	28.84	11.92 - 69.76	<0.001	0.17	0.12 - 0.26	<0.001
Platelet count † (* 10 ⁹ /L)	<100	6.88	3.61 - 13.13	<0.001	0.16	0.05 - 0.49	0.001
	≥ 100	1.00			1.00		
SOFA score †		1.63	1.54 - 1.74	<0.001	0.62	0.57 - 0.67	<0.001
D-dimer(mg/L) \$	<0.5	1.00			1.00		
	0.5 - 1.0	1.53	0.63 - 3.67	0.346	0.82	0.64 - 1.06	0.129
	>1.0	3.34	1.55 - 7.19	0.002	0.57	0.45 - 0.73	<0.001
Ferritin(ng/mL) \$	<500	1.00			1.00		
	≥500	4.11	2.28 - 7.41	<0.001	0.52	0.42 - 0.64	<0.001
CRP\$ (mg/L)		1.0003	0.999 - 1.001	0.080	0.99	0.99 - 1.00	0.360
LDH\$ (IU/L)	<450	1.00			1.00		
	≥ 450	4.88	2.72 - 8.75	<0.001	0.53	0.43 - 0.66	<0.001
PaO2/FiO2‡	<100 (severe)	25.64	14.8 - 44.41	<0.001	6.5e-08	4.3e-08 - 9.9e-08	<0.001
	100-200(moderate)	5.97	3.05 - 11.69	<0.001	0.19	0.10 - 0.36	<0.001
	>200 (Mild)	1.00			1.00		
Interval from onset of symptoms to admission		1.05	0.96 - 1.14	0.334	0.97	0.94 - 1.00	0.058
Vasopressor support		11.36	7.79 - 16.56	<0.001	0.03	0.004 - 0.22	0.001
Invasive ventilation support		19.57	12.21 - 31.35	<0.001	0.01	0.002 - 0.09	<0.001

† Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: \$ inflammatory biomarkers values were measured at day 0, 3 and day 7.SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

PLACID TRIAL Mortality Assessment

498

499 Table 3: Multivariable Fine and Gray Model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Multivariable Analysis (Model A)						Multivariable Analysis (Model B)					
		Mortality			Discharged alive			Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00			1.00			1.00		
	41-59	1.23	0.60 - 2.49	0.572	1.03	0.81 - 1.32	0.825	1.55	0.65 - 3.71	0.325	0.95	0.74 - 1.21	0.671
	≥60	1.44	0.67 - 3.09	0.347	0.94	0.70 - 1.26	0.675	1.72	0.67 - 4.46	0.261	0.78	0.57 - 1.06	0.110
Comorbidities	No Comorbidities	1.00			1.00			1.00			1.00		
	1	1.20	0.69 - 2.10	0.515	0.91	0.72 - 1.14	0.407	1.31	0.59 - 2.91	0.509	0.87	0.67 - 1.11	0.254
	2 or More	1.76	1.02 - 3.03	0.041	0.89	0.69 - 1.14	0.329	2.68	1.26 - 5.70	0.011	0.69	0.52 - 0.93	0.014
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00								
	5-10	3.34	1.21 - 9.19	0.020	0.81	0.64 - 1.03	0.095						
	>10	9.97	3.65 - 27.13	<0.001	0.39	0.26 - 0.58	<0.001						
SOFA score †		1.22	1.11 - 1.35	<0.001	0.75	0.68 - 0.83	<0.001						
D-dimer(mg/L) §	<0.5							1.00					
	0.5 - 1.0							1.29	0.54 - 3.10	0.565	0.84	0.65 - 1.09	0.198
	>1.0							2.50	1.14 - 5.48	0.022	0.64	0.49 - 0.82	<0.001
Ferritin(ng/mL) §	<500							1.00			1.00		
	≥500							2.67	1.44 - 4.96	0.002	0.69	0.55 - 0.86	0.001
LDH§ (IU/L)	<450							1.00			1.00		
	≥ 450							2.96	1.60 - 5.45	0.001	0.68	0.55 - 0.85	0.001
PaO2/FiO2†	<100 (severe)	3.47	1.64 - 7.37	0.001	3.1e-07	1.7e-07 - 5.7e-07	<0.001						
	100-200(moderate)	1.91	0.91 - 4.004	0.087	0.401	0.19 - 0.85	0.016						
	>200 (Mild)	1.00			1.00								

500

501 †Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: § Inflammatory biomarkers values were measured at day 0, 3 and day 7

502 (Model A) Multivariable Fine and Gray model for Age, comorbidities with Laboratory Parameters, (Model B) Multivariable Fine and Gray model for Age, comorbidities with
503 inflammatory biomarker values

504 SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

505

506

PLACID TRIAL Mortality Assessment

Figure 1:





Figure 1: Line graph for SOFA score showing difference between survivors and non survivors

Figure showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2•30 and 3•05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

PLACID TRIAL Mortality Assessment

Supplementary

Flowchart for the study protocol



The Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID) Trial recruited 464 eligible patients for the study. The primary outcome at 28 days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomisation and two patients did not receive the intervention after randomisation as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients out of whom 386 survived and 65 died.

BMJ Open

Factors associated with mortality among moderate and severe COVID 19 patients in India: A secondary analysis of a Randomised Controlled Trial

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2021-050571.R3
Article Type:	Original research
Date Submitted by the Author:	25-Aug-2021
Complete List of Authors:	<p>Mammen, Joy; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Kumar, Snehil; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Thomas, Lovely; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Kumar, Gunjan ; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Zachariah, Anand; Christian Medical College and Hospital Vellore, Medicine</p> <p>Jeyaseelan, Lakshmanan; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Peter, John Victor; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Agarwal, Anup; ICMR, Clinical Trial and Health Systems Research Unit</p> <p>Mukherjee, Aparna; ICMR, Clinical Trials and Health Systems Research Unit</p> <p>Chatterjee, Pranab; ICMR, Translational Global Health Policy and Research Cell</p> <p>Bhatnagar, Tarun ; National Institute of Epidemiology, ICMR School of Public Health</p> <p>Rasalam, Jess; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Chacko, Binila; Christian Medical College and Hospital Vellore, Medical Intensive Care Unit</p> <p>Mani, Thenmozhi; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Joy, Melvin; Christian Medical College and Hospital Vellore, Biostatistics</p> <p>Rupali, Priscilla; Christian Medical College and Hospital Vellore, Infectious Diseases</p> <p>Murugesan, Malathi; Christian Medical College and Hospital Vellore, Hospital Infection Control Committee</p> <p>Daniel, Dolly; Christian Medical College and Hospital Vellore, Transfusion Medicine and Immunohaematology</p> <p>Latha, B; Madras Medical College, Transfusion Medicine</p> <p>Bundas, Sunita; SMS Medical College and Hospital, Transfusion Medicine</p> <p>Kumar, Vivek; Sir HN Reliance Foundation Hospital and Research Centre, Critical Care</p> <p>Dosi, Ravi; Sri Aurobindo Institute of Medical Sciences, Respiratory Medicine</p>

	Khambholja, Janakkumar; Smt NHL Municipal Medical College, Internal Medicine de Souza, Rosemarie; BYL Nair Charitable Hospital, Department of Medicine Chander, B; Gandhi Medical College and Hospital, Internal Medicine Bahadur, Shalini; Government Institute of Medical Sciences, Pathology Dube, Simmi; Gandhi Medical College Bhopal, Internal Medicine Suri, Amit; Atal Bihari Vajpayee Institute of Medical Sciences and Ram Manohar Lohia Hospital, Pulmonary Medicine Jindal, Aikaj ; Satguru Pratap Singh Hospitals, Transfusion Medicine Shrivastav, Om ; Kasturba Hospital for Infectious Diseases, Infectious diseases Barge, Vijay; RSCM Government Medical College, Medicine Bajpayee, Archana; AIIMS Jodhpur, Transfusion Medicine Malhotra, Pankaj; Post Graduate Institute of Medical Education and Research, Department of Internal Medicine Singh, Neha; AIIMS Patna, Transfusion Medicine Tambe, Muralidhar; Department of Community Medicine, B J Govt. Medical College Sharma, Nimisha; ESIC Medical College and Hospital Faridabad, Transfusion Medicine Bhat, Shreepad; Smt Kashibai Navale Medical College and General Hospital, Internal Medicine Kaulgud, Ram; Karnataka Institute of Medical Sciences, Internal Medicine Gurtoo, Anil; Lady Hardinge Medical College, Internal Medicine Reddy, Himanshu; King George Medical College, Internal Medicine Upadhyay, Kamlesh; Byramjee Jeejabhoy Medical College, Internal medicine Jain, Ashish ; Mahatma Gandhi Medical College and Hospital, Respiratory Medicine Patel, Tinkal; Government Medical College, Internal Medicine Nagori, Irfan; GMERS Medical College Gotri Vadodara, Medicine Jha, Pramod ; Sumandeep Vidyapeeth University, Internal Medicine Babu, KV; Sri Venkateswara Institute of Medical Sciences, Transfusion Medicine Aparna, C; Kurnool Medical College, Pathology Panjwani, Sunil; Government Medical College, Internal Medicine Natarajan, M; Madurai Medical College, Internal Medicine Baldi, Milind; Mahatama Gandhi Memorial Medical College, Internal Medicine Khadke, Vrushali; Poona Hospital and Research Centre, Interventional Pulmonology Dua, Seema; Super Speciality Paediatric Hospital and Teaching Hospital, Transfusion Medicine Singh, Ravindraa; Aditya Birla Memorial Hospital, Transfusion Medicine Sharma, Ashish; R. D. Gardi Medical College, Medicine Sharma, Jayashree; Seth GS Medical College and KEM Hospital, Transfusion Medicine Gokhale, Yojana; Lokmanya Tilak Municipal Medical College and General Hospital, Internal Medicine Yadav, Pragya; National Institute of Virology, Maximum Containment Laboratory, ICMR Sapkal, Gajanan; National Institute of Virology, Diagnostic Virology Group, ICMR Kaushal, Himanshu; National Institute of Virology, Human Influenza Group, ICMR Kumar, Saravana; National Institute of Epidemiology, Epidemiology and Biostatistics Division
Primary Subject	Public health

Heading:	
Secondary Subject Heading:	Public health
Keywords:	COVID-19, Public health < INFECTIOUS DISEASES, Adult intensive & critical care < INTENSIVE & CRITICAL CARE

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

PLACID TRIAL Mortality Assessment

**Factors associated with mortality among moderate and severe COVID 19 patients in India:
A secondary analysis of a Randomised Controlled Trial**

Joy J Mammen, Snehil Kumar, Lovely Thomas, Gunjan Kumar, Anand Zachariah, Lakshmanan Jeyaseelan, John Victor Peter, Anup Agarwal, Aparna Mukherjee, Pranab Chatterjee, Tarun Bhatnagar, Jess Elizabeth Rasalam, Binila Chacko, Thenmozhi Mani, Melvin Joy, Priscilla Rupali, Malathi Murugesan, Dolly Daniel, B Latha , Sunita Bundas, Vivek Kumar, Ravi Dosi, Janakkumar R Khambholja, Rosemarie de Souza, B Thrilok Chander, Shalini Bahadur, Simmi Dube, Amit Suri, Aikaj Jindal, Om Shrivastav, Vijay Barge, Archana Bajpayee, Pankaj Malhotra, Neha Singh, Muralidhar Tambe, Nimisha Sharma, Shreepad Bhat, Ram S Kaulgud, Anil Gurtoo, D Himanshu Reddy, Kamlesh Upadhyay, Ashish Jain, Tinkal C Patel, Irfan Nagori, Pramod R Jha, KV Sreedhar Babu, C Aparna, Sunil Jodharam Panjwani, M Natarajan, Milind Baldi, Vrushali Khirid Khadke, Seema Dua, Ravindraa Singh, Ashish Sharma, Jayashree Sharma, Yojana A Gokhale, Pragya D Yadav, Gajanan Sapkal, Himanshu Kaushal, V Saravana Kumar

Corresponding Author:

Joy J Mammen MD

Professor & Head

Address

Department of Transfusion Medicine and Immunohaematology

Christian Medical College Vellore

Tamil Nadu, India

Pin code 632004

E mail joymammen@cmcvellore.ac.in

Phone +91 416 228 2536

Fax +91 416 223 2035

PLACID TRIAL Mortality Assessment

32

Name	Designation and affiliation	Full Address
Joy J Mammen	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Snehil Kumar	Senior Resident, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lovely Thomas	Assistant Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Gunjan Kumar	Scientist C, Clinical Trial & Health Systems Research Unit, Indian Council of Medical Research (ICMR)	New Delhi, India.
Anand Zachariah	Professor, Department of Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Lakshmanan Jeyaseelan	Professor, Clinical Epidemiology Unit, Christian Medical College	Vellore, Tamil Nadu, India.
John Victor Peter	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Anup Agarwal	Consultant, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Aparna Mukherjee	Scientist E, Clinical Trial & Health Systems Research Unit, ICMR	New Delhi, India.
Pranab Chatterjee	Scientist B, Translational Global Health Policy and Research Cell, ICMR	New Delhi, India.
Tarun Bhatnagar	Scientist E, ICMR School of Public Health, ICMR -National Institute of Epidemiology	Chennai, Tamil, India
Jess Elizabeth Rasalam	Fellow, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Binila Chacko	Professor, Department of Critical Care Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
Thenmozhi Mani	Senior Demonstrator, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Melvin Joy	Associate Research Officer, Department of Biostatistics, Christian Medical College	Vellore, Tamil Nadu, India.
Priscilla Rupali	Professor, Department of Infectious Diseases, Christian Medical College	Vellore, Tamil Nadu, India.
Malathi Murugesan	Assistant Professor, Hospital Infection Control Committee, Christian Medical College	Vellore, Tamil Nadu, India.
Dolly Daniel	Professor, Department of Transfusion Medicine, Christian Medical College	Vellore, Tamil Nadu, India.
B Latha	Professor, Department of Transfusion Medicine, Madras Medical College	Chennai, Tamil Nadu, India
Sunita Bundas	Professor, Department of Immunohaematology and Transfusion Medicine, SMS Medical College and Hospital	Jaipur, Rajasthan, India
Vivek Kumar	Professor, Department of Critical Care Medicine, Sir H N Reliance Foundation Hospital	Mumbai, Maharashtra, India
Ravi Dosi	Professor, Department of Respiratory Medicine, Sri Aurobindo Institute of Medical Sciences	Indore, Madhya Pradesh, India
Janakkumar R Khambholja	Professor, Department of Internal Medicine, Smt NHL Municipal Medical College	Ahmedabad, Gujarat, India
Rosemarie de Souza	Professor, Department of Internal Medicine, BYL Nair Charitable Hospital & TN Medical College	Mumbai, Maharashtra, India
B Thrilok Chander	Professor, Department of Internal Medicine, Gandhi Medical College, Secunderabad	Secunderabad, Telangana, India
Shalini Bahadur	Associate Professor, Department of Pathology, Government Institute of Medical Sciences	Greater Noida, Uttar Pradesh, India
Simmi Dube	Professor, Department of Internal Medicine, Gandhi Medical College	Bhopal, Madhya Pradesh, India
Amit Suri	Professor, Department of Pulmonary Medicine, Atal Bihari Vajpayee Institute of Medical Sciences and RML Hospital	New Delhi, India
Aikaj Jindal	Professor, Department of Transfusion Medicine, Satguru Partap Singh Hospitals	Ludhiana, Punjab, India
Om Shrivastav	Professor, Department of Infectious Diseases, Kasturba Hospital	Mumbai, Maharashtra, India
Vijay Barge	Professor, Department of Medicine, RSCM Government Medical College	Kolhapur, Maharashtra, India
Archana Bajpayee	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Jodhpur, Rajasthan, India
Pankaj Malhotra	Professor, Department of Internal Medicine, Post Graduate Institute of Medical Education and Research	Chandigarh, India
Neha Singh	Professor, Department of Transfusion Medicine, All India Institute of Medical Sciences	Patna, Bihar, India
Muralidhar Tambe	Professor, Department of Community Medicine Byramjee Jeejeebhoy Medical College	Pune, Maharashtra, India

PLACID TRIAL Mortality Assessment

Nimisha Sharma	Professor, Department of Immunohematology and blood transfusion, ESIC Medical College and Hospital	Faridabad, Haryana, India
Shreepad Bhat	Professor, Department of Internal Medicine, Smt. Kashi Bai Navale Medical College	Pune, Maharashtra, India
Ram S Kaulgud	Professor, Department of Internal Medicine, Karnataka Institute of Medical Sciences	Hubali, Karnataka, India
Anil Gurtoo	Professor, Department of Internal Medicine, Lady Hardinge Medical College	New Delhi, India
D Himanshu Reddy	Professor, Department of Internal Medicine, King George's Medical University	Lucknow, Uttar Pradesh, India
Kamlesh Upadhyay	Professor, Department of Internal Medicine, Byramjee Jeejeebhoy Medical College	Ahmedabad, Gujarat, India
Ashish Jain	Professor, Department of Respiratory Medicine, Mahatma Gandhi University of Medical Sciences and Technology	Jaipur, Rajasthan, India
Tinkal C Patel	Professor Department of Internal Medicine, Government Medical College	Surat, Gujarat, India
Irfan Nagori	Professor, Department of Medicine, GMERS Medical College and Hospital	Vadodara, Gujarat, India
Pramod R Jha	Professor, Department of Internal Medicine, Sumandeep, Vidyapeeth and Dhiraj Hospital	Vadodara, Gujarat, India
KV Sreedhar Babu	Professor, Department of Transfusion Medicine, Sri Venkateswra Institute of Medical Sciences	Tirupati, Andhra Pradesh, India
C Aparna	Professor, Department of Pathology, Kurnool Medical College	Kurnool, Andhra Pradesh, India
Sunil Jodharam Panjwani	Professor, Department of Internal Medicine, Government Medical College	Bhavnagar, Gujarat, India
M Natarajan	Professor, Department of Internal Medicine, Madurai Medical College	Madurai, Tamil Nadu, India
Milind Baldi	Professor, Department of Respiratory Medicine, Mahatma Gandhi Memorial Medical College	Indore, Madhya Pradesh, India
Vrushali Khirid Khadke	Professor, Department of Interventional Pulmonology, Poona Hospital and Research Center	Pune, Maharashtra, India
Seema Dua	Professor, Department of Transfusion Medicine, Super Specialty Paediatric Hospital and Post Graduate Teaching Institute	Noida, Uttar Pradesh, India
Ravindraa Singh	Professor, Department of Transfusion Medicine, Aditya Birla Memorial Hospital	Pune, Maharashtra, India
Ashish Sharma	Professor, Department of Medicine, RD Gardi Medical College	Ujjain, Madhya Pradesh, India
Jayashree Sharma	Professor, Department Transfusion Medicine, Seth GS Medical College and KEM Hospital	Mumbai, Maharashtra, India
Yojana A Gokhale	Professor, Department of Internal Medicine, Lokmanya Tilak Municipal Medical College & General Hospital	Mumbai, Maharashtra, India
Pragya D Yadav	Scientist E, Maximum Containment Laboratory, ICMR – National Institute Virology	Pune, Maharashtra, India
Gajanan Sapkal	Scientist E, Diagnostic Virology Group, ICMR – National Institute Virology	Pune, Maharashtra, India
Himanshu Kaushal	Scientist B, Human Influenza group, ICMR – National Institute Virology	Pune, Maharashtra, India
V Saravana Kumar	Scientist B, Epidemiology and Biostatistics Division, ICMR – National Institute of Epidemiology	Chennai, Tamil Nadu, India

33

PLACID TRIAL Mortality Assessment

Abstract

Objective:

Large data on the clinical characteristics and outcome of COVID-19 in the Indian population is scarce. We analysed the factors associated with mortality in a cohort of moderate and severely ill COVID-19 patients enrolled in a randomised trial on convalescent plasma.

Design:

Secondary analysis of data from a Phase II, Open Label, Randomised Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID TRIAL).

Setting:

39 public and private hospitals across India, during the study period 22 April 2020 to 14 July 2020.

Participants:

Of the 464 patients recruited, two were lost to follow-up, nine withdrew consent and two patients did not receive the intervention after randomisation. The cohort of 451 participants with known outcome at 28-days was analysed.

Primary outcome measure:

Factors associated with all-cause mortality at 28-days post-enrolment.

Results:

The mean (SD) age was 51±12.4 years; 76.7% were males. Admission SOFA score was 2.4±1.1. Non-invasive ventilation, invasive ventilation and vasopressor therapy were required in 98.9%, 8.4% and 4.0% respectively. The 28-day mortality was 14.4%. Median time from symptom onset to hospital admission was similar in survivors (4 days; IQR 3-7) and non survivors (4 days; IQR 3-6). Patients with two or more co-morbidities had 2.25 (95% CI: 1.18-4.29, p=0.014) times risk of death. When compared with survivors, admission IL-6 levels were higher (p<0.001) in non-survivors and increased further on Day 3. On multivariable Fine and Gray model, severity of illness (sub-distribution hazard ratio (SHR) 1.22, 95% CI: 1.11-1.35, p<0.001), PaO₂/FiO₂ ratio <100 (3.47, 1.64-7.37, p=0.001), Neutrophil Lymphocyte ratio (NLR) >10 (9.97, 3.65-27.13, p<0.001), D-dimer >1.0 mg/L (2.50, 1.14-5.48, p=0.022), ferritin ≥500 ng/mL (2.67, 1.44-4.96, p=0.002) and LDH ≥450 IU/L (2.96, 1.60-5.45, p=0.001) were significantly associated with death.

Conclusion:

In this cohort of moderate and severely ill COVID-19 patients, severity of illness, underlying co-morbidities and elevated levels of inflammatory markers were significantly associated with death.

PLACID TRIAL Mortality Assessment

67

Trial Registration:

CTRI/2020/04/024775

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

PLACID TRIAL Mortality Assessment

90

91 **Article Summary**

92 **Strengths and limitations of this study**

93 **Strengths**

94 There is no study from India, with representation from multiple states that has detailed the

95 clinical profile and evaluated for factors associated with death. This study may help with strategic

96 planning at a national level.

97 The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28,

98 did not differ across the trial arms, therefore the present analysis need not be adjusted for

99 convalescent plasma intervention.

100 There may be variability of treatment provided in the multiple centres, however, care was taken

101 that patients received best standard of care for COVID-19 dictated by the best available evidence at

102 the time and guidelines for the management of COVID-19 issued by health authorities of the Indian

103 government.

104 **Limitations**

105 The laboratory and biomarker assays for ferritin, lactate dehydrogenase, C reactive protein, and D-

106 dimer were conducted using tests from different manufacturers.

107 Participants of this study may not comprise a true observational cohort as this was a *post hoc*

108 analysis of a randomised control trial data. Our study did not analyse the effect of SARS-CoV-2

109 variants causing a high mortality in younger population during the second wave of COVID-19

110 infection, and therefore extrapolation to the general population must be carefully qualified.

111

112

113

114

PLACID TRIAL Mortality Assessment**Introduction**

The first human case of Corona Virus Disease 19 (COVID-19) caused by the novel coronavirus (named Severe Acute Respiratory Syndrome Coronavirus 2, SARS-CoV-2) was reported in Wuhan City, China in December 2019. On 30 January 2020, the World Health Organisation (WHO) declared that the outbreak of COVID-19 constituted a Public Health Emergency of International Concern (1). Based on the high level of global spread and the severity of COVID-19, on 11 March 2020, the Director-General of the WHO declared the COVID-19 outbreak a pandemic (2). The sudden outbreak followed by rapid spread in a globalised world, resulted in a huge burden on the healthcare system, besides affecting the socio-economic well-being among all nations.

In India, the disease was first detected on 30 January 2020 in the state of Kerala, in a student who returned from Wuhan (3,4). After a brief, initial respite, the virus spread at a rapid pace in India, resulting in more than 10 million confirmed cases as of December, 2020 with more than 145,000 deaths (5).

Most patients diagnosed with COVID-19 experience mild to moderate respiratory illness, fever, dry cough and fatigue and recover without requiring special treatment (6). Oxygen desaturation is the hallmark of progression. Patients with underlying medical problems like cardiovascular disease, diabetes, chronic respiratory disease, and cancer are more likely to develop serious illness. These patients may develop viral pneumonia, with resultant dyspnoea and hypoxaemia which may progress to respiratory or multi- system failure and even death (7). There is paucity of large-scale data on the clinical characteristics, outcomes of COVID-19 in the Indian population and evaluation of risk factors with an unfavourable outcome at a national level. Identification of such potential risk factors is important to anticipate medical treatment and to reduce the mortality burden for severe COVID-19 illness by proactive interventions.

The Indian Council of Medical Research (ICMR) conducted a randomised trial (A Phase II, Open Label, Randomised Controlled Trial to Assess the Safety and Efficacy of Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease, PLACID TRIAL) to determine the effectiveness and safety of convalescent plasma in patients with moderate and severely ill COVID-19 to limit progression of disease (8). Patients received standard of care for COVID-19 in keeping with the institutional protocols, based on the best available evidence at the time and guidelines for the management of COVID-19 issued by national health authorities. Participants in the intervention arm received two doses of 200 mL of convalescent plasma, transfused 24 hours apart, in addition

PLACID TRIAL Mortality Assessment

to standard of care. The control arm did not receive any additional therapy. The study concluded that the use of convalescent plasma was not associated with a reduction in 28-day mortality (8). The aim of this analysis was to identify risk factors associated with mortality by mining the data collected from the cohort enrolled in the PLACID TRIAL (8).

Methods

Participants

The study enrolled patients from 39 different hospitals, of which, 29 were teaching public hospitals and 10 were private facilities across 14 states and union territories. Patients over the age of 18 years who were confirmed to have COVID-19 based on a positive SARS-CoV-2 RT-PCR test and presenting with moderate and severely illness with either a partial pressure of oxygen in arterial blood/fraction of inspired oxygen (PaO₂/FiO₂) ratio between 200-300 or respiratory rate >24/min and decreased oxygen saturation on room air (SpO₂ < 93%) were included during the study period from 22 April to 14 July 2020. As per the guidelines issued by the Ministry of Health, Government of India at the time of conduct of the study, the subset of patients with the above criteria but with a respiratory rate between 24 and 30/min were classified as moderate disease. Those with respiratory rate >30 breaths/min were classified as severe disease (9). Patients were followed up for 28-days and assessed for their health status and all-cause mortality. Ethics approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) as well as from the Institutional Review Boards (IRB) / Institutional Ethics Committees of all the participating hospitals. Written consent was obtained from the patients or their families before enrolling in the study.

Data

Data was obtained from the ICMR PLACID TRIAL database collected in structured paper case record forms and entered in Research Electronic Data Capture system (REDCap, version 8.5 Vanderbilt University, TN). The trial protocol was registered with the Clinical Trial Registry of India (CTRI/2020/04/024775). After trial completion, based on co-operative agreement between the centres, and IRB permission, the data was shared and analysed further, to explore for other meaningful results. No separate ethical clearance was taken for this study.

PLACID TRIAL Mortality Assessment

Demographic, clinical, laboratory tests and outcome data were collected prospectively. Clinical symptoms, need for organ support (respiratory, renal, haemodynamic), laboratory tests (complete blood count, coagulation profile, serum biochemical profile, renal and liver function tests) were monitored serially on day of enrolment (day 0) and on days 1, 3, 5, 7, 14 and 28. Inflammatory biomarkers [lactate dehydrogenase (LDH), serum ferritin, and C-reactive protein (CRP)] were tested at admission and on day 3 and 7 whereas Interleukin 6 (IL-6) was done at admission and on day 3.

The outcome of interest was all-cause day 28 mortality. In addition, we looked for association between laboratory parameters and mortality.

Statistical Methods:

Mean and standard deviation (SD) or median and inter-quartile range (IQR) were used for continuous variables as appropriate, and for categorical variables, number and proportions were used. To find the association between mortality and study variables, Chi-square test/Fisher's exact test were used. To find the mean difference across the groups, independent t-test was used. Similarly, Mann Whitney U test was used to compare median difference. The primary end-point was all-cause mortality (event of interest) at day 28 from the time of enrolment, discharged alive (competing event) and hospital admission after day 28 (censored), whichever was earlier. Discharged alive was treated as a competing event because the event of "discharged alive" precludes the event of all-cause mortality. The variables that were statistically significant or clinically important were considered in the multivariable Fine and Gray regression model. However, if a variable was expected to have collinear concern or had sparse data, it was not included in the analysis. Two multivariable models were developed. The first model included clinical and laboratory parameters tested on day 0, 1, 3, 5, 7, 14 and 28 while the second included inflammatory biomarkers tested on day 0, 3 and 7, after adjusting for age and comorbidities. Variables that were considered included parameters that were strongly associated with mortality at univariate analysis or those known from previous literature to be strongly associated with outcome. For certain laboratory markers such as D-dimer, ferritin and LDH, clinically relevant thresholds were used for the analysis rather than using these data as continuous variables. The clinically relevant thresholds for these variables were set as >1.0 mg/L for D-dimer, ≥ 500 mg/mL for Ferritin and ≥ 450 IU/L for LDH. The threshold for Ferritin of $500 \mu\text{g/L}$ was based on the cut-off value for the diagnosis of Hemophagocytic lymphohistiocytosis (HLH) as well as some preliminary evidence in COVID that a

PLACID TRIAL Mortality Assessment

threshold of >500 µg/L was associated with invasive ventilator dependence (10). Similarly, traditionally a threshold of <0.5 mg/L is used to exclude pulmonary thromboembolism; in this context two thresholds were used, 0.5 to 1.0 mg/L and >1.0 mg/L (11). The model assumption was verified using log-log S (t) plots and Global test. A p-value of less than 0.05 levels was considered as statistically significant. All statistical analysis were performed using STATA version 16.0 (StataCorp. 2019. College Station, TX).

Patient and public involvement

Patients and the public were not involved in the design, conduct, reporting, or dissemination plans of our research. The study results will be disseminated to the study participants via their treating doctors.

Results

The PLACID Trial recruited 464 eligible patients for the study. The primary outcome at 28-days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomisation and two patients did not receive the intervention after randomisation as a matched donor was not available. The cohort with known outcome at 28 days thus comprised 451 patients (*supplementary*).

The primary outcome of the PLACID TRIAL, disease progression or all-cause mortality at day 28, did not differ across the trial arms, therefore the analysis did not adjust for convalescent plasma intervention. The distribution of patients in the intervention and control arms were 50.3% (n=227) and 49.7% (n=224), respectively. The mean (SD) age of the cohort was 51 ± 12.4 years; 76.7% were males. Table 1 shows the distribution of demographic variables and clinical parameters in the study population.

The most common presenting symptoms were shortness of breath (91.6%), fatigue (78.7%), cough (68.5%) and fever (35%). Comorbidities were present in 59.9% of patients; 31.7% had any one comorbidity and 28.2% had two or more comorbidities. The most frequent comorbidities were diabetes (43.5%), hypertension (37.5%), obesity (6.9%) and Chronic Obstructive Pulmonary Disease (COPD) (3.3%). There was history of smoking in 8.2%. The time from onset of symptoms to admission was four days (IQR 3-7 days). Majority of the patients required non-invasive (98.9%) ventilatory support. The median duration of respiratory support was six days (IQR 4-10 days). In

PLACID TRIAL Mortality Assessment

this cohort, 4% patients required vasopressor support. None of the patients required Extra Corporeal Membrane Oxygenation (ECMO) or dialysis support.

The all-cause mortality at 28-days was 14.4% (95%CI: 11.5-17.9, n=65). Median time from symptom onset to hospital admission was four days in survivors (IQR 3-7 days) and non-survivors (IQR 3-6 days). The frequency of shortness of breath, cough and fatigue were similar in survivors and non-survivors; however, the presence of fever at admission was significantly ($p=0.042$) associated with death (table 1). Other than COPD and CKD (chronic kidney disease), other comorbidities were not significantly associated with death (table 1). Admission Sequential Organ Failure Assessment (SOFA) score was higher in non survivors. The need for invasive mechanical ventilation, duration of invasive mechanical ventilation and vasopressor therapy were associated with death (table 1).

On univariate analysis (table 2), there was an association between increasing age and mortality. Patients with two or more comorbidities had a 2.25 (95%CI: 1.18-4.29, $p=0.014$) times increased chance of mortality. There was a strong mortality association for platelet count $<100 \times 10^9/L$ (SHR 6.88, 95%CI: 3.61-13.13, $p<0.001$), neutrophil lymphocyte ratio (NLR) >10 (28.84, 11.92-69.76, $p<0.001$), LDH ≥ 450 IU/L (4.88, 2.72-8.75, $p<0.001$), D-dimer >1 mg/L (3.34, 1.55-7.19, $p=0.002$) and ferritin ≥ 500 ng/mL (4.11, 2.28-7.41, $p<0.001$). Admission IL-6 levels were significantly ($p<0.001$) higher (76.00, 18.27-171.77) in non-survivors than in survivors (18.51, 4.26-56.86). By day 3, IL-6 levels dropped to 11.6 (2.64-45.84) in survivors while it nearly doubled in non-survivors (140.35, 21.56-427.36). CRP did not show any statistical significance (1.0003, 0.999-1.001, $p=0.080$).

The need for invasive ventilation and vasopressors were associated with death (table 2). Increasing SOFA score was associated with mortality (1.63, 1.54-1.74, $p<0.001$). The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score progressively increased between the two groups over time (figure 1). Mortality also proportionately increased with lower PaO_2/FiO_2 values with sub-distribution hazard ratio (SHR) of 25.64 (14.8-44.41, $p<0.001$) in the severe group as compared to the mild group.

Two models were run for multivariable Fine and Gray regression model over a period of time (table 3). Model A included age, comorbidities, PaO_2/FiO_2 , NLR and SOFA score. Model A revealed significant sub-distribution hazard ratios for PaO_2/FiO_2 ratio <100 (3.47, 1.64-7.37, $p=0.001$), NLR >10 (9.97, 3.65-27.13, $p<0.001$), SOFA score (1.22, 1.11- 1.35, $p<0.001$) after adjusting for age and

PLACID TRIAL Mortality Assessment

comorbidities. Model B included age, comorbidities, D-dimer, ferritin and LDH. D-dimer >1 mg/L (2.50, 1.14-5.48, p=0.022), ferritin ≥500 ng/mL (2.67, 1.44-4.96, p=0.002) and LDH ≥450 IU/L (2.96, 1.60-5.45, p=0.001), were associated with mortality after adjusting for age and comorbidities (table 3). IL-6 was omitted from the model as it was not measured on Day 7.

Discussion

In this study that enrolled patients in the PLACID trial from across India, SOFA score and clinical biomarkers like D-dimer, LDH and ferritin were identified as factors that could predict increased risk of death in moderate and severely ill COVID-19 patients. The definition of clinical grading of severity is different in India as compared to other countries (12–16). Mortality of critically ill COVID-19 patients varies significantly among already published case series and ranges from 16% to 78% (17–23). Two studies from Wuhan, which included moderate as well as critically ill patients, showed mortality rates of 3.77% and 14.14% (24,25). This wide variability can be explained by differences in the age of the population, distribution of risk factors, health system responses, varied treatment protocols and disparate follow-up times. In a series of critically ill patients in China, the 28-day ICU mortality was 61.5% (26). In a multicentric study from Italy, the mortality risk for patients without respiratory failure at admission was 1% after 15 days, while survival in patients with moderate-to-severe respiratory failure (PaO₂/FiO₂ ≤200 mm Hg) at admission was only 56% at 15 days (27). The fatality rate reported in Europe and the United States of America was significantly higher than in China (28). Therefore, findings obtained in a specific country might not be automatically extrapolated and national cohorts must be studied.

In our study population, mortality increased with age; this pattern was observed in other countries affected by COVID-19. Age seemed to affect the time from hospitalisation to death. Age-specific death rates were quite similar in studies from Asia, Europe and North America (29). In South Korea, Italy, France, Germany, England and Wales, and Spain, the COVID-19 attributed mortality rates rose by about 12% per year whereas the United States and Wuhan, China had a lower rate of increase of about 9.5% per year of age (30). In a meta-analysis of 611,583 subjects, the overall mortality was 12.10%; the lowest mortality rate was reported from China (3.1%) and the highest in the United Kingdom (20.8%) and New York State (20.99%). Among the patients included in the meta-analysis, 23.2% were ≥80 years of age; mortality was highest in these patients. The largest increase in

PLACID TRIAL Mortality Assessment

mortality risk was observed in patients aged 60 to 69 years as compared with those aged 50 to 59 years (odds ratio 3.13, 95% CI: 2.61-3.76) (31).

The presence of comorbidities significantly increases the death risk due to COVID-19. A higher risk of mortality was seen in our patients who had CKD and COPD. A meta-analysis, including 1389 COVID-19 patients, with 19.7% having severe disease showed a significant association of CKD with severe COVID-19 with pooled odds ratio as 3.03 (32). Similarly, the estimated mortality risk in patients with COPD was three times than those without ($p < 0.05$) (33). We found that 43.5% of our patients had diabetes which is markedly higher when compared with patients from Korea which showed that 16.97% had diabetes mellitus (34). Our analysis showed that the presence of diabetes was not significantly different between survivors and non-survivors (42.5% vs. 49.2%, $p = 0.310$), in contrast to the study from South Korea (34) which showed a much higher mortality among diabetic patients than in those without (20.0% vs. 4.8%). Hypertension and obesity were not significantly different among survivors and non-survivors in our study. However, the presence of two or more comorbidities was associated with mortality in our study.

The Fine-Gray model identified prognostic markers for mortality, most notably age ≥ 60 years, $\text{PaO}_2/\text{FiO}_2$ ratio < 100 , NLR > 10 , platelet count $< 100 \times 10^9/\text{L}$, ferritin $> 500\text{ng/mL}$, LDH $> 450\text{ IU/L}$ and D-dimer $> 1\text{mg/L}$. Our study findings were similar when compared with studies from Wuhan (35). Older age, leukocytosis, and high LDH level have been reported to be risk factors associated with in-hospital death in other studies also (36–38). IL-6 levels were significantly different in survivors and non-survivors at admission. By day 3 survivors had reducing IL-6 while it nearly doubled in non-survivors.

Mortality was higher amongst patients requiring invasive mechanical ventilation (SHR 19.57, 12.21-31.35, $p < 0.001$) and those requiring vasopressors (SHR 11.36, 7.79-16.56, $p < 0.001$). However, the median duration of invasive ventilation for survivors was 12 days (IQR 2, 14) and that for non-survivors was one day (IQR 1, 3). These results suggest that the sickest patients probably die very early in the course of hospitalisation, while patients with acute respiratory failure requiring ventilatory support may survive with prolonged ventilatory support. Therefore, invasive ventilation should be offered in a timely manner and effectively provided.

In our study, the SOFA score was recognised as a valuable tool that could be used to prognosticate outcome of patients with COVID-19. Competing risk regression models showed that the increase in

PLACID TRIAL Mortality Assessment

SOFA score was related to mortality, with a clearly divergent pattern between the two groups. Thus, an increasing SOFA score over time may be a factor that can be used to identify a subset of patients who may have an unfavourable outcome. Studies have shown that the SOFA score could be used to evaluate severity and 60-day mortality of COVID-19 with the optimal cut-off score of 5 (39).

The limitations of this study include, the variability of treatment provided in the multiple centres. The participants of this study may not comprise a true observational cohort, as this was a *post hoc* analysis of a randomised control trial data and extrapolation to the general population must be carefully qualified. Our study did not analyse the effect of SARS-CoV-2 variants causing a high mortality in younger population during the second wave of COVID-19 infection and this may limit generalisability of the data to the second wave. Despite these limitations, this study provides a comprehensive overview of prognostic factors in moderate and severely ill COVID-19 patients that included patients from across the country.

Conclusion

Older age, multiple comorbidities, low PaO₂/FiO₂ ratio and elevated levels of inflammatory markers are associated with worse prognosis. Serial SOFA score can be used for prognostication. Understanding the symptoms, burden of comorbidities and systematic monitoring of key laboratory parameters offer opportunities for targeted intervention in COVID-19 with the use of anti-inflammatory or immunomodulatory agents.

Figure legend

Figure 1 showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2•30 and 3•05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

FOOTNOTES:

Authors Contributions:

Study design: JJM, LJ, JVP

PLACID TRIAL Mortality Assessment

Clinical Management: SK, LT, AZ, JER, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG.

Data collection: JJM, SK, LT, AZ, AA, AM, GK, PC, TB, JER, PR, MM, BC, BL, SUB, VK, RD, JRK, RDS, BTC, SB, SD, ASU, AJJ, OS, VB, AB, PM, NS, MT, NMS, SPB, RSK, AG, DHR, KU, AJ, TCP, IN, PRJ, KVS, CA, SJP, MN, MB, VKK, SMD, RVS, AS, JS, YAG, PDY, GS, HK, VSK.

Data Analysis: JJM, SK, LJ, TM, MJ, PR, MM, DD, JVP.

Data Interpretation: JJM, SK, AZ, LJ, JER, TM, MJ, DD, JVP

Manuscript writing: JJM, SK, LT, AZ, LJ, JER, BC, TM, MJ, PR, MM, DD, JVP, AA, AM, GK, HK, PC, TB

Study Administration: JJM, BC, JVP, AA, AM, GK, HK, PC, TB

Funding support: The funding source for the primary study was ICMR (RFC No. ECD/NTF/20/2020-21/Covid Dated 23.07.2020). However, they had no financial interest in the investigational product. No other author has any competing financial or non-financial interest.

Competing interests: 'None declared'. All authors have completed the ICMJE uniform disclosure form at www.icmje.org.

Ethical Approval: Ethical approval was obtained from the ICMR Central Ethics Committee on Human Research (CECHR-002/2020) based in the National Center for Disease Informatics and Research, Indian Council of Medical Research, Bengaluru, Karnataka, as well as from the Institutional Review Boards (IRB) /Institutional Ethics Committees of all the participating hospitals.

Data availability statement: Data will be made available, upon request, and must be accompanied by a brief proposal outlining the analysis plan. A signed data access agreement might be needed to ensure data safety and compliance with national rules about data sharing.

Acknowledgement: We acknowledge ICMR for providing us the data collected during PLACID trial which was analysed in the present study.

References:

1. Mullen L, Potter C, Gostin LO, Cicero A, Nuzzo JB. An analysis of International Health Regulations Emergency Committees and Public Health Emergency of International Concern Designations. BMJ Glob Health. 2020 Jun 1;5(6):e002502.

PLACID TRIAL Mortality Assessment

- 386 2. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Bio-Medica Atenei Parm.* 2020 19;91(1):157–60.
- 388 3. Perappadan BS. India's first coronavirus infection confirmed in Kerala. *The Hindu* [Internet]. 2020 Jan 30 [cited 2020 Nov 17]; Available from: <https://www.thehindu.com/news/national/indias-first-coronavirus-infection-confirmed-in-kerala/article30691004.ece>
- 391 4. Andrews MA, Areekal B, Rajesh KR, Krishnan J, Suryakala R, Krishnan B, et al. First confirmed case of COVID-19 infection in India: A case report. *Indian J Med Res.* 2020 May 1;151(5):490.
- 393 5. MoHFW | Home [Internet]. [cited 2021 Jan 3]. Available from: <https://www.mohfw.gov.in/>
- 394 6. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol Orlando Fla.* 2020 Jun;215:108427.
- 396 7. Wu C, Chen X, Cai Y, Xia J 'an, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med.* 2020 01;180(7):934–43.
- 399 8. Agarwal A, Mukherjee A, Kumar G, Chatterjee P, Bhatnagar T, Malhotra P. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre randomised controlled trial (PLACID Trial). *BMJ* [Internet]. 2020 Oct 22 [cited 2020 Nov 16];371. Available from: <https://www.bmj.com/content/371/bmj.m3939>
- 403 9. Government of India, Ministry of Health and Family Welfare, Directorate General of Health Services. CLINICAL MANAGEMENT PROTOCOL FOR COVID-19-Version 3. Government of India; 2020.
- 406 10. Qeadan F, Tingey B, Gu LY, Packard AH, Erdei E, Saeed AI. Prognostic Values of Serum Ferritin and D-Dimer Trajectory in Patients with COVID-19. *Viruses.* 2021 05;13(3).
- 408 11. Kearon C, de Wit K, Parpia S, Schulman S, Afilalo M, Hirsch A, et al. Diagnosis of Pulmonary Embolism with d-Dimer Adjusted to Clinical Probability. *N Engl J Med.* 2019 Nov 28;381(22):2125–34.
- 411 12. Varghese GM, John R, Manesh A, Karthik R, Abraham OC. Clinical management of COVID-19. *Indian J Med Res.* 2020 May 1;151(5):401.
- 413 13. Clinical management of COVID-19 [Internet]. [cited 2021 Jan 3]. Available from: <https://www.who.int/publications/i/item/clinical-management-of-covid-19>
- 415 14. Zhang S-Y, Lian J-S, Hu J-H, Zhang X-L, Lu Y-F, Cai H, et al. Clinical characteristics of different subtypes and risk factors for the severity of illness in patients with COVID-19 in Zhejiang, China. *Infect Dis Poverty.* 2020 Jul 8;9(1):85.
- 418 15. Xia L, Chen J, Friedemann T, Yang Z, Ling Y, Liu X, et al. The Course of Mild and Moderate COVID-19 Infections—The Unexpected Long-Lasting Challenge. *Open Forum Infect Dis* [Internet]. 2020 Sep 1;7(ofaa286). Available from: <https://doi.org/10.1093/ofid/ofaa286>
- 421 16. Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai N, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis.* 2020 Jun 1;20(6):669–77.

PLACID TRIAL Mortality Assessment

- 423 17. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al. Risk Factors Associated
424 With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy. *JAMA*
425 *Intern Med.* 2020 01;180(10):1345–55.
- 426 18. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019
427 novel coronavirus in Wuhan, China. *Lancet Lond Engl.* 2020 15;395(10223):497–506.
- 428 19. Wang Y, Lu X, Li Y, Chen H, Chen T, Su N, et al. Clinical Course and Outcomes of 344 Intensive
429 Care Patients with COVID-19. *Am J Respir Crit Care Med.* 2020 Apr 8;201(11):1430–4.
- 430 20. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized
431 Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA.* 2020 Mar
432 17;323(11):1061–9.
- 433 21. Bhatraju PK, Ghassemieh BJ, Nichols M, Kim R, Jerome KR, Nalla AK, et al. Covid-19 in Critically Ill
434 Patients in the Seattle Region - Case Series. *N Engl J Med.* 2020 21;382(21):2012–22.
- 435 22. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and Outcomes
436 of 21 Critically Ill Patients With COVID-19 in Washington State. *JAMA.* 2020 28;323(16):1612–4.
- 437 23. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult
438 inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet Lond Engl.* 2020
439 28;395(10229):1054–62.
- 440 24. Zhang J, Wang X, Jia X, Li J, Hu K, Chen G, et al. Risk factors for disease severity, unimprovement,
441 and mortality in COVID-19 patients in Wuhan, China. *Clin Microbiol Infect.* 2020 Jun 1;26(6):767–
442 72.
- 443 25. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased
444 patients with coronavirus disease 2019: retrospective study. *BMJ.* 2020 Mar 26;368:m1091.
- 445 26. Yang X, Yu Y, Xu J, Shu H, Xia J 'an, Liu H, et al. Clinical course and outcomes of critically ill
446 patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective,
447 observational study. *Lancet Respir Med.* 2020 May 1;8(5):475–81.
- 448 27. Santus P, Radovanovic D, Saderi L, Marino P, Cogliati C, De Filippis G, et al. Severity of respiratory
449 failure at admission and in-hospital mortality in patients with COVID-19: a prospective
450 observational multicentre study. *BMJ Open [Internet].* 2020 Oct 10;10(10). Available from:
451 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7549463/>
- 452 28. Yamamoto N, Bauer G. Apparent difference in fatalities between Central Europe and East Asia
453 due to SARS-COV-2 and COVID-19: Four hypotheses for possible explanation. *Med Hypotheses.*
454 2020 Nov;144:110160.
- 455 29. Demographics of COVID-19 Deaths [Internet]. Ined - Institut national d'études démographiques.
456 [cited 2020 Nov 16]. Available from: <https://dc-covid.site.ined.fr/en/>
- 457 30. Goldstein JR, Lee RD. Demographic perspectives on the mortality of COVID-19 and other
458 epidemics. *Proc Natl Acad Sci U S A.* 2020 Sep 8;117(36):22035–41.
- 459 31. Bonanad C, García-Blas S, Tarazona-Santabalbina F, Sanchis J, Bertomeu-González V, Fácila L, et
460 al. The Effect of Age on Mortality in Patients With COVID-19: A Meta-Analysis With 611,583
461 Subjects. *J Am Med Dir Assoc.* 2020 Jul 1;21(7):915–8.

PLACID TRIAL Mortality Assessment

32. Henry BM, Lippi G. Chronic kidney disease is associated with severe coronavirus disease 2019 (COVID-19) infection. *Int Urol Nephrol*. 2020 Mar 28;1–2.
33. Venkata VS, Kiernan G. COVID-19 AND COPD: POOLED ANALYSIS OF OBSERVATIONAL STUDIES. *CHEST*. 2020 Oct 1;158(4):A2469.
34. D A, K L, Ds L, Ys L, Ss M. Mortality Rate and Predictors of Mortality in Hospitalized COVID-19 Patients with Diabetes. *Healthc Basel Switz* [Internet]. 2020 Sep 13 [cited 2020 Nov 16];8(3). Available from: <https://europepmc.org/article/med/32933191>
35. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* [Internet]. 2020 May 7;55(5). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144257/>
36. Li M, Cheng B, Zeng W, Chen S, Tu M, Wu M, et al. Analysis of the Risk Factors for Mortality in Adult COVID-19 Patients in Wuhan: A Multicenter Study. *Front Med* [Internet]. 2020 [cited 2020 Nov 16];7. Available from: <https://www.frontiersin.org/articles/10.3389/fmed.2020.00545/full>
37. Albitar O, Ballouze R, Ooi JP, Sheikh Ghadzi SM. Risk factors for mortality among COVID-19 patients. *Diabetes Res Clin Pract*. 2020 Aug 1;166:108293.
38. Holman N, Knighton P, Kar P, O’Keefe J, Curley M, Weaver A, et al. Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: a population-based cohort study. *Lancet Diabetes Endocrinol*. 2020 Oct 1;8(10):823–33.
39. Zheng Yang, Qinming Hu, Fei Huang, Shouxin Xiong, Yi Sun. The prognostic value of the SOFA score in patients with COVID-19: a retrospective, observational study. 2020 Oct 26 [cited 2020 Nov 20]; Available from: <https://europepmc.org/article/ppr/ppr231113>

PLACID TRIAL Mortality Assessment

Table 1: Distribution of demographic variables and clinical parameters in enrolled patients and comparison of survivors and non-survivors in the cohort

Variables	Overall (n=451) N (%)	Survivor (n=386) N (%)	Non survivor (n=65) N (%)	P value
Age (Mean ± SD)	51±12.4	50±12.4	56±11.3	<0.001*
Age				
≤ 40	104 (23.1)	97 (25.1)	7 (10.8)	0.004
41-59	225 (49.9)	194 (50.3)	31 (47.7)	
≥60	122 (27.1)	95 (24.6)	27 (41.5)	
Gender: Male	346 (76.7)	294(76.2)	52 (80.0)	0.499
Blood group				0.518
A	104(23.1)	91(23.6)	13(20.0)	
B	164(36.4)	140(36.3)	24(36.9)	
AB	25(5.5)	19(4.9)	6(9.2)	
O	158(35.0)	136(35.2)	22(33.8)	
History of smoking	37(8.2)	32(8.3)	5(7.7)	0.866
Comorbidities and Chronic illness				
Diabetes	196 (43.5)	164 (42.5)	32 (49.2)	0.310
Hypertension	169 (37.5)	139 (36.0)	30 (46.2)	0.118
Chronic obstructive pulmonary disease	15 (3.3)	10 (2.6)	5 (7.7)	0.050
Obesity ≥ 30	31 (6.9)	25 (6.5)	6 (9.2)	0.426
Chronic kidney disease	17 (3.8)	11 (2.8)	6 (9.2)	0.024
Coronary artery disease	31 (6.9)	23 (6.0)	8 (12.3)	0.106
Cerebrovascular disease	4 (0.9)	3 (0.8)	1 (1.5)	0.465
Symptoms at admission				
Shortness of breath	413 (91.6)	351 (90.9)	62 (95.4)	0.232
Fever	158 (35.0)	128 (33.2)	30 (46.2)	0.042
Cough	309 (68.5)	259 (67.1)	50 (76.9)	0.115
Fatigue	354 (78.7)	301 (78.2)	53 (81.5)	0.541
Severity of illness score				
SOFA score at admission*	2.40 ± 1.06	2.30±0.93	3.05±1.49	<0.001
Treatment				
Vasopressor	18 (4.0)	1 (0.3)	17 (26.6)	<0.001
Non-Invasive Ventilation (NIV)	446 (98.9)	383 (99.2)	63 (96.9)	0.101
Invasive ventilation	38 (8.4)	4 (1.04)	34 (52.31)	<0.001
Interval between symptoms onset to admission ‡	4 (3, 7)	4 (3, 7)	4 (3, 6)	0.996
Duration of respiratory support days ‡	6 (4, 10)	6 (4, 9.5)	6 (3, 10)	0.689
Duration of invasive ventilation days ‡	1 (1,3)	12 (2, 14)	1 (1, 3)	0.020

PLACID TRIAL Mortality Assessment

Duration of hospital stay	14 (10, 18)	14 (11, 19)	8 (5, 14)	<0.001
days ‡				
499				
500	‡ Median (IQR) days in days – Mann Whitney U test was used			
501	*Mean ± SD – Independent t test was used			

For peer review only

PLACID TRIAL Mortality Assessment

Table 2: Univariate Fine and Gray model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Univariate Analysis					
		Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00		
	41-59	2.04	0.90 - 4.66	0.089	0.80	0.64 - 1.01	0.057
	≥60	3.51	1.53 - 8.07	0.003	0.56	0.42 - 0.73	<0.001
Gender	Male	1.19	0.64 - 2.19	0.582	0.87	0.70 - 1.09	0.228
Blood Group	O	1.00			1.00		
	A	0.94	0.48 - 1.87	0.866	0.89	0.69 - 1.15	0.389
	B	1.13	0.63 - 2.01	0.689	0.93	0.75 - 1.18	0.578
	AB	2.01	0.80 - 5.05	0.139	0.66	0.39 - 1.11	0.116
Comorbidities	No Comorbidities	1.00			1.00		
	1	1.62	0.82 - 3.21	0.166	0.79	0.63 - 0.99	0.044
	2 or More	2.25	1.18 - 4.29	0.014	0.70	0.55 - 0.88	0.003
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00		
	5-10	4.90	1.80 - 13.32	0.002	0.72	0.56 - 0.93	0.013
	>10	28.84	11.92 - 69.76	<0.001	0.17	0.12 - 0.26	<0.001
Platelet count † (* 10 ⁹ /L)	<100	6.88	3.61 - 13.13	<0.001	0.16	0.05 - 0.49	0.001
	≥ 100	1.00			1.00		
SOFA score †		1.63	1.54 - 1.74	<0.001	0.62	0.57 - 0.67	<0.001
D-dimer(mg/L) \$	<0.5	1.00			1.00		
	0.5 - 1.0	1.53	0.63 - 3.67	0.346	0.82	0.64 - 1.06	0.129
	>1.0	3.34	1.55 - 7.19	0.002	0.57	0.45 - 0.73	<0.001
Ferritin(ng/mL) \$	<500	1.00			1.00		
	≥500	4.11	2.28 - 7.41	<0.001	0.52	0.42 - 0.64	<0.001
CRP ^s (mg/L)		1.0003	0.999 - 1.001	0.080	0.99	0.99 - 1.00	0.360
LDH ^s (IU/L)	<450	1.00			1.00		
	≥ 450	4.88	2.72 - 8.75	<0.001	0.53	0.43 - 0.66	<0.001
PaO ₂ /FiO ₂ ‡	<100 (severe)	25.64	14.8 - 44.41	<0.001	6.5e-08	4.3e-08 - 9.9e-08	<0.001
	100-200(moderate)	5.97	3.05 - 11.69	<0.001	0.19	0.10 - 0.36	<0.001
	>200 (Mild)	1.00			1.00		
Interval from onset of symptoms to admission		1.05	0.96 - 1.14	0.334	0.97	0.94 - 1.00	0.058
Vasopressor support		11.36	7.79 - 16.56	<0.001	0.03	0.004 - 0.22	0.001
Invasive ventilation support		19.57	12.21 - 31.35	<0.001	0.01	0.002 - 0.09	<0.001

† Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: \$ inflammatory biomarkers values were measured at day 0, 3 and day 7.SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

PLACID TRIAL Mortality Assessment

502

503 Table 3: Multivariable Fine and Gray Model for baseline characteristics, laboratory parameters and inflammatory biomarkers

Variables		Multivariable Analysis (Model A)						Multivariable Analysis (Model B)					
		Mortality			Discharged alive			Mortality			Discharged alive		
		SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value	SHR	95% CI	P value
Age	≤40	1.00			1.00			1.00			1.00		
	41-59	1.23	0.60 - 2.49	0.572	1.03	0.81 - 1.32	0.825	1.55	0.65 - 3.71	0.325	0.95	0.74 - 1.21	0.671
	≥60	1.44	0.67 - 3.09	0.347	0.94	0.70 - 1.26	0.675	1.72	0.67 - 4.46	0.261	0.78	0.57 - 1.06	0.110
Comorbidities	No Comorbidities	1.00			1.00			1.00			1.00		
	1	1.20	0.69 - 2.10	0.515	0.91	0.72 - 1.14	0.407	1.31	0.59 - 2.91	0.509	0.87	0.67 - 1.11	0.254
	2 or More	1.76	1.02 - 3.03	0.041	0.89	0.69 - 1.14	0.329	2.68	1.26 - 5.70	0.011	0.69	0.52 - 0.93	0.014
Neutrophil/Lymphocyte ratio †	<5	1.00			1.00								
	5-10	3.34	1.21 - 9.19	0.020	0.81	0.64 - 1.03	0.095						
	>10	9.97	3.65 - 27.13	<0.001	0.39	0.26 - 0.58	<0.001						
SOFA score †		1.22	1.11 - 1.35	<0.001	0.75	0.68 - 0.83	<0.001						
D-dimer(mg/L) §	<0.5							1.00					
	0.5 - 1.0							1.29	0.54 - 3.10	0.565	0.84	0.65 - 1.09	0.198
	>1.0							2.50	1.14 - 5.48	0.022	0.64	0.49 - 0.82	<0.001
Ferritin(ng/mL) §	<500							1.00			1.00		
	≥500							2.67	1.44 - 4.96	0.002	0.69	0.55 - 0.86	0.001
LDH§ (IU/L)	<450							1.00			1.00		
	≥ 450							2.96	1.60 - 5.45	0.001	0.68	0.55 - 0.85	0.001
PaO2/FiO2‡	<100 (severe)	3.47	1.64 - 7.37	0.001	3.1e-07	1.7e-07 - 5.7e-07	<0.001						
	100-200(moderate)	1.91	0.91 - 4.004	0.087	0.401	0.19 - 0.85	0.016						
	>200 (Mild)	1.00			1.00								

504

505 †Laboratory Parameters were measured at day 0, 1,3,5,7 and day 14: § Inflammatory biomarkers values were measured at day 0, 3 and day 7

506 (Model A) Multivariable Fine and Gray model for Age, comorbidities with Laboratory Parameters, (Model B) Multivariable Fine and Gray model for Age, comorbidities with
507 inflammatory biomarker values

508 SHR: Sub-distribution Hazard ratio; CI: Confidence Interval

509

510

PLACID TRIAL Mortality Assessment

Figure 1:





Figure 1: Line graph for SOFA score showing difference between survivors and non survivors

Figure showing serial Sequential Organ Failure Assessment (SOFA) score among survivors and non-survivors. Increasing SOFA score was associated with mortality. The mean SOFA score at day 0 was 2.30 and 3.05 for survivors and non-survivors respectively. The difference in the SOFA score showed divergence between the two groups over time.

PLACID TRIAL Mortality Assessment

Supplementary

Flowchart for the study protocol



The Convalescent Plasma to Limit COVID-19 Associated Complications in Moderate Disease (PLACID) Trial recruited 464 eligible patients for the study. The primary outcome at 28 days was not available for two patients who were lost to follow-up after discharge; nine patients withdrew consent after randomisation and two patients did not receive the intervention after randomisation as a matched donor was not available. The cohort with known outcome at 28 days thus comprised of 451 patients out of whom 386 survived and 65 died.